



Canadian Academy of Child and Adolescent Psychiatry
Académie canadienne de psychiatrie de l'enfant et de l'adolescent

Implications of Cannabis Legalization on Youth and Young Adults

Phil Tibbo, MD, FRCPC, FCPA¹; Candice E. Crocker, MSc, PhD²; Raymond W. Lam, MD, FRCPC³;
Jeff Meyer, MD, PhD, FRCPC⁴; Jitender Sareen, MD, FRCPC⁵;
Katherine J. Aitchison, BA (Hons), MA (Oxon), BM BCh, PhD, FRCPsych⁶

A position statement developed by the Canadian Psychiatric Association's Research Committee and approved by the CPA's Board of Directors on February 17, 2017.

The Canadian Psychiatric Association (CPA) is the national leading authority on psychiatric matters in Canada with a stated vision of a **strong profession for a healthy population**. As an evidence-based profession, part of the CPA's mission is to advocate for policies that allow the best possible mental wellness for Canadians, working collaboratively with governments and mental health stakeholders.

The federal government has committed to legalizing and regulating access to cannabis in 2017. The government appreciates the complexity of this process as evidenced by their writing a discussion paper and forming a Task Force on Marijuana Legalization and Regulation (Task Force) that was charged with engaging multi-stakeholder groups and knowledge leaders to better inform the design of the new framework.

This CPA position statement is focused on youth and young adults as mental illness and substance use disorders often begin in this age group, with the objective to ensure optimal mental health outcomes. The CPA recognizes the Canadian Medical Association's (CMA) submission to the Task Force that covers recommendations related to overall health as well as mental health.

Cannabis is the most commonly used illicit drug for Canadian youth, and Canadian youth are the top users of cannabis in the developed world. While there are variations by jurisdictions, approximately 22% of youth aged 15 to 19 reported using cannabis during the past year; in this group, 20% report daily or almost daily use.

The CPA recognizes the following evidence-based knowledge:

¹ Professor and Dr Paul Janssen Chair in Psychotic Disorders, Department of Psychiatry, Dalhousie University

² Department of Psychiatry, Dalhousie University and Department of Diagnostic Imaging, Nova Scotia Health Authority

³ Professor and BC Leadership Chair in Depression Research, Department of Psychiatry, University of British Columbia

⁴ Professor and Canada Research Chair in Neurochemistry of Depression, Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health and Department of Psychiatry, University of Toronto

⁵ Professor and Head, Department of Psychiatry, University of Manitoba

⁶ Professor, Department of Psychiatry and Adjunct Professor, Department of Medical Genetics, University of Alberta

© Copyright 2017, Canadian Psychiatric Association. This document may not be reproduced without written permission of the CPA. Members' comments are welcome. Please address all comments and feedback to: President, Canadian Psychiatric Association, 141 Laurier Avenue West, Suite 701, Ottawa, ON K1P 5J3; Tel: 613-234-2815; Fax: 613-234-9857; e-mail: president@cpa-apc.org. Reference 2017-34s.

Note: It is the policy of the Canadian Psychiatric Association to review each position paper, policy statement and clinical practice guideline every five years after publication or last review. Any such document that has been published more than five years ago and does not explicitly state it has been reviewed and retained as an official document of the CPA, either with revisions or as originally published, should be considered as a historical reference document only.

- The human brain undergoes a maturational process during adolescence that includes reorganization, refinements and functional improvements. This is driven by changes in brain grey matter (GM) due to synaptic pruning (elimination of underutilized or unnecessary neural connections), and white matter (WM) due to myelination.
- These changes continue at least until the mid-20s and thus brain maturation is vulnerable during this time to stressors/insults.
- The endocannabinoid system plays a role in this brain maturation and thus exogenous cannabinoids from cannabis can affect this process directly in a negative way.
- Regular cannabis use in youth and young adults can affect aspects of cognition, including attention, memory, processing speed, visuospatial functioning and overall intelligence. Worse performance is related to earlier adolescent onset of use. Abstinence following regular use may improve some, but not all, of these cognitive domains.
- Early and regular use increases the risk of developing a primary psychotic illness in those individuals who are vulnerable. Vulnerability factors are not currently clear, but may include factors such as childhood trauma and genetics. In those young adults who have developed psychosis, continued cannabis use worsens long-term symptom and functional outcomes.
- Cannabis may increase the risk of depression and early regular use is associated with younger age of onset of symptoms of psychosis and of bipolar disorder.
- Cannabis with high tetrahydrocannabinol (THC) content (high potency) can result in significantly worse mental health and cognitive outcomes, including worsening of panic disorder and other anxiety disorders.
- Early age of use of cannabis increases the potential for adult dependence to cannabis.
- Cannabis may be associated with increased progression to other illicit drug use in the context of particular factors (e.g., high frequency and early age of use).

As this CPA position statement is focusing on youth and young adults, it recognizes the possibility of cannabis use in young women of reproductive age. Prenatal cannabis exposure (PCE) may have adverse effects on cognitive development, behaviour and academic achievement in offspring due to the effects of cannabis on the fetal

endocannabinoid system modulating neurodevelopment. PCE has been hypothesized to potentially compromise the brain so that a “second hit” (postnatal stressors/ environmental triggers) precipitates the emergence of these abnormalities.

The CPA acknowledges and agrees with the CMA recommendations to the Task Force. With respect to protection of mental wellness for youth and young adults the CPA highlights the following:

- Since regular cannabis use is associated with increased risk of schizophrenia, and may also negatively interact with depression, bipolar and anxiety disorders due to its biological effects on brain maturation, and since mental disorders frequently start before the age of 25, age of access to cannabis should not be prior to age 21, with restrictions on quantity and THC potency for those between 21 and 25 years of age.
- Significant support is needed for public health education and resources targeting youth and young adults, in collaboration with mental health stakeholder groups.
- Significant support of further biological and psychosocial research is needed to better understand the impact of cannabis and its legalization on mental health.
- Expand support for prevention, early identification and cannabis cessation treatments (i.e., using change-based treatment models including harm reduction strategies) within the framework of mental health and addictions.
- Prudent consideration of advertising and marketing guidelines with clear markings of THC and cannabidiol content, as well as consistent public health warning messaging, including regarding potential adverse consequences of the use of cannabis during pregnancy, are needed.

References

1. Adamson P. Child well-being in rich countries: a comparative overview. Innocenti Report Card 11. Florence (Italy): United Nations International Children’s Emergency Fund Office of Research; 2013.
2. Andréasson S, Allebeck P, Engström A, et al. Cannabis and schizophrenia: a longitudinal study of Swedish conscripts. *Lancet*. 1987;2(8574):1483–1486.
3. Alpar A, Tortoriello G, Calvigioni D, et al. Endocannabinoids modulate cortical development by configuring Slit2/Robo1 signalling. *Nat Commun*. 2014;5:4421.
4. Alpar A, Di Marzo V, Harkany T. At the tip of an iceberg: Prenatal marijuana and its possible relation to neuropsychiatric outcome in the offspring. *Biol Psychiatry*. 2016;79(7):e33–45.

5. Arseneault L, Cannon M, Poulton R, et al. Cannabis use in adolescence and risk for adult psychosis: longitudinal prospective study. *BMJ*. 2002;325(7374):1212–1213.
6. Behnke M, Smith VC, Committee on Substance Abuse, et al. Prenatal substance abuse: short- and long-term effects on the exposed fetus. *Pediatrics*. 2013;131(3):e1009–1024.
7. Benes FM, Turtle M, Khan Y, et al. Myelination of a key relay zone in the hippocampal formation occurs in the human brain during childhood, adolescence, and adulthood. *Arch Gen Psychiatry*. 1994;51(6):477–484.
8. Bossong MG, Niesink RJ. Adolescent brain maturation, the endogenous cannabinoid system and the neurobiology of cannabis-induced schizophrenia. *Prog Neurobiol*. 2010;92(3):370–385.
9. Bray S. Age-associated patterns in gray matter volume, cerebral perfusion and BOLD oscillations in children and adolescents. *Hum Brain Mapp*. 2017;Epub 2017 Jan 25.
10. Canadian Medical Association (CMA). Legalization, regulation and restriction of access to marijuana. CMA submission to the Government of Canada Task Force on Marijuana Legalization and Regulation [Internet]. Ottawa (ON): CMA; 2016 [cited 2016 Aug 29]. Available from: <https://www.cma.ca/Assets/assets-library/document/en/advocacy/submissions/2016-aug-29-cma-submission-legalization-and-regulation-of-marijuana-e.pdf>.
11. Camchong J, Lim KO, Kumra S. Adverse effects of cannabis on adolescent brain development: A longitudinal study. *Cereb Cortex*. 2016;Epub 2016 Feb 26.
12. Carr JA, Norman RM, Manchanda R. Substance misuse over the first 18 months of specialized intervention for first episode psychosis. *Early Interv Psychiatry*. 2009;3(3):221–225.
13. Degenhardt L, Stockings E, Patton G, et al. The increasing global health priority of substance use in young people. *Lancet Psychiatry*. 2016;3(3):251–264.
14. Duff G, Argaw A, Cecyre B, et al. Cannabinoid receptor CB2 modulates axon guidance. *PLoS One*. 2013;8(8):e70849.
15. Fergusson DM, Boden JM, Horwood LJ. Psychosocial sequelae of cannabis use and implications for policy: findings from the Christchurch Health and Development Study. *Soc Psychiatry Psychiatr Epidemiol*. 2015;50(9):1317–1326.
16. Fergusson DM, Horwood LJ, Swain-Campbell N. Cannabis use and psychosocial adjustment in adolescence and young adulthood. *Addiction*. 2002;97(9):1123–1135.
17. Gibbs M, Winsper C, Marwaha S, et al. Cannabis use and mania symptoms: a systematic review and meta-analysis. *J Affect Disord*. 2015;171:39–47.
18. Giedd JN, Blumenthal J, Jeffries NO, et al. Brain development during childhood and adolescence: a longitudinal MRI study. *Nat Neurosci*. 1999;2(10):861–863.
19. Glass M, Dragunow M, Faull RL. Cannabinoid receptors in the human brain: a detailed anatomical and quantitative autoradiographic study in the fetal, neonatal and adult human brain. *Neuroscience*. 1997;77(2):299–318.
20. Health Canada. Canadian tobacco, alcohol and drugs survey [Internet]. Ottawa (ON): Health Canada, 2013 [cited 2017 Jan 10]. Available from: <http://healthycanadians.gc.ca/science-research-sciences-recherches/data-donnees/ctads-ectad/summary-sommaire-2013-eng.php>.
21. Henquet C, Krabbendam L, Spauwen J, et al. Prospective cohort study of cannabis use, predisposition for psychosis, and psychotic symptoms in young people. *BMJ*. 2005;330(7481):11.
22. Horwood LJ, Fergusson DM, Coffey C, et al. Cannabis and depression: an integrative data analysis of four Australasian cohorts. *Drug Alcohol Depend*. 2012;126(3):369–378.
23. Kedzior KK, Laeber LT. A positive association between anxiety disorders and cannabis use or cannabis use disorders in the general population—a meta-analysis of 31 studies. *BMC Psychiatry*. 2014;14:136.
24. Kelly C, Castellanos FX, Tomaselli O, et al. Distinct effects of childhood ADHD and cannabis use on brain functional architecture in young adults. *Neuroimage Clin*. 2017;13:188–200.
25. Large M, Sharma S, Compton MT, et al. Cannabis use and earlier onset of psychosis: a systematic meta-analysis. *Arch Gen Psychiatry*. 2011;68(6):555–561.
26. Lev-Ran S, Roerecke M, Le Foll B, et al. The association between cannabis use and depression: a systematic review and meta-analysis of longitudinal studies. *Psychol Med*. 2014;44(4):797–810.
27. McGrath J, Welham J, Scott J, et al. Association between cannabis use and psychosis-related outcomes using sibling pair analysis in a cohort of young adults. *Arch Gen Psychiatry*. 2010;67(5):440–447.
28. National Academies of Sciences, Engineering, and Medicine. The health effects of cannabis and cannabinoids: The current state of evidence and recommendations for research. Washington (DC): The National Academies Press, 2017.
29. Peters A. The effects of normal aging on myelin and nerve fibers: a review. *J Neurocytol*. 2002;31(8–9):581–593.
30. Molina-Holgado E, Vela JM, Arévalo-Martin A, et al. Cannabinoids promote oligodendrocyte progenitor survival: involvement of cannabinoid receptors and phosphatidylinositol-3 kinase/Akt signaling. *J Neurosci*. 2002;22(22):9742–9753.
31. Pfefferbaum A, Mathalon DH, Sullivan EV, et al. A quantitative magnetic resonance imaging study of changes in brain morphology from infancy to late adulthood. *Arch Neurol*. 1994;51(9):874–887.
32. Renard J, Krebs MO, Le Pen G, et al. Long-term consequences of adolescent cannabinoid exposure in adult psychopathology. *Front Neurosci*. 2014;8:361.
33. Richardson KA, Hester AK, McLemore GL. Prenatal cannabis exposure—The “first hit” to the endocannabinoid system. *Neurotoxicol Teratol*. 2016;58:5–14.
34. Richmond S, Johnson KA, Seal ML, et al. Development of brain networks and relevance of environmental and genetic factors: A systematic review. *Neurosci Biobehav Rev*. 2016;71:215–239.
35. Secades-Villa R, Garcia-Rodriguez O, Jin CJ, et al. Probability and predictors of the cannabis gateway effect: a national study. *Int J Drug Policy*. 2015;26(2):135–142.
36. Silveri MM, Dager AD, Cohen-Gilbert JE, et al. Neurological signatures associated with alcohol and drug use in the human adolescent brain. *Neurosci Biobehav Rev*. 2016;70:244–259.
37. Smith AM, Mioduszewski O, Hatchard T, et al. Prenatal marijuana exposure impacts executive functioning into young adulthood: an fMRI study. *Neurotoxicol Teratol*. 2016;58:53–59.
38. Squeglia LM, Gray KM. Alcohol and drug use and the developing brain. *Curr Psychiatry Rep*. 2016;18(5):46.
39. Tzilos GK, Reddy MK, Caviness CM, et al. Getting higher: co-occurring drug use among marijuana-using emerging adults. *J Addict Dis*. 2014;33(3):202–209.
40. Zvolensky MJ, Cogle JR, Johnson KA, et al. Marijuana use and panic psychopathology among a representative sample of adults. *Exp Clin Psychopharmacol*. 2010;18(2):129–134.

