

Oncogenic *ras*-induced downregulation of a pro-apoptotic protease caspase-2 is required for malignant transformation of intestinal epithelial cells.

Byong Hoon Yoo[‡], Yanfei Wang[‡], Mete Erdogan[‡], Takehiko Sasazuki[§], Senji Shirasawa[¶], Laurent Corcos^{*}, Kanaga Sabapathy^{||} and Kirill V. Rosen[‡]

From [‡]Departments of Pediatrics & Biochemistry and Molecular Biology, Atlantic Research Centre, Dalhousie University, Halifax, Nova Scotia, Canada, [§]Department of Pathology, Research Institute, International Medical Center of Japan, Tokyo, Japan, [¶]Department of Cell Biology, School of Medicine, Fukuoka University, Fukuoka, Japan, ^{*} INSERM, Faculté de Médecine, Brest, France, ^{||} Laboratory of Molecular Carcinogenesis, National Cancer Centre, Singapore

Address correspondence to: Kirill Rosen, Atlantic Research Centre, Room C-302, CRC, Dalhousie University, 5849 University Avenue, Halifax, Nova Scotia, Canada B3H 4H7. Tel.: (902) 494-7088. Fax: (902) 494-1394. email: kirill.rosen@dal.ca

Running title: *ras* transforms cells by downregulating caspase-2

Resistance of carcinoma cells to anoikis, apoptosis that is normally induced by loss of cell-to-extracellular matrix (ECM) adhesion, is thought to be essential for the ability of these cells to form primary tumors, invade adjacent tissues and metastasize to distant organs. The current knowledge about the mechanisms by which cancer cells evade anoikis is far from complete. In an effort to understand these mechanisms we found that *ras*, a major oncogene, downregulates protease caspase-2 (that initiates certain steps of the cellular apoptotic program) in malignant human and rat intestinal epithelial cells. This downregulation could be reversed by inhibition of a protein kinase Mek, a mediator of Ras signaling. We also found that enforced downregulation of caspase-2 in non-malignant intestinal epithelial cells by RNA interference protected them from anoikis. Furthermore, the reversal of the effect of Ras on caspase-2 achieved by the expression of exogenous caspase-2 in detached *ras*-transformed intestinal epithelial cells promoted well established apoptotic events, such as the release of the pro-apoptotic mitochondrial factors cytochrome c and HtrA2/Omi into the cytoplasm of these cells, significantly enhanced their anoikis susceptibility and blocked their long-term growth in the absence of adhesion to the ECM. Finally, the blockade of the effect of Ras on caspase-2 substantially suppressed growth of tumors formed by the *ras*-transformed cells in mice. We conclude that *ras*-induced downregulation of caspase-2 represents a novel mechanism by which oncogenic Ras protects

malignant intestinal epithelial cells from anoikis, promotes their anchorage-independent growth and allows them to form tumors *in vivo*.

Many normal epithelia are organized *in vivo* into cellular monolayers, which are attached to the form of the extracellular matrix (ECM) referred to as basement membrane (BM). Detachment of epithelial cells from the ECM causes their apoptotic death (1;2), a phenomenon termed anoikis (2). Unlike normal epithelia, carcinomas (cancers derived from epithelial cells) typically represent three-dimensional disorganized multicellular masses in which cell-ECM contacts are significantly changed. It is known in this regard that carcinoma cells typically grow as multilayers and at least some of these cells are detached from the BM. It is also well established that cancer cells often produce BM-degrading enzymes and this allows tumors to invade adjacent tissues (3). Furthermore, at advanced stages of cancer cellular aggregates detach from the primary tumor and seed in other organs where they give rise to metastases (4;5). However, even though carcinoma cells are deprived of normal contacts with the BM during tumor progression, many of these cells do not undergo anoikis (4;5).

Several lines of evidence support the notion that anoikis resistance represents a critical prerequisite for carcinoma progression. First, cancer cells can typically survive and grow being detached from the ECM as colonies in soft agar. This property represents one of the most stringent criteria for malignant transformation that are presently being used (6;7). Second, we and others established that activation of oncoproteins, such as Ras (1), EGF receptor (8) and β -catenin (9) or loss

of tumor suppressor genes, such as PTEN (10), can block anoikis of cancer cells. Furthermore, we and others found that treatments that reverse anoikis resistance of tumor cells also suppress their ability to form primary tumors (11-15) and metastases (5;11;14;16;17). In addition we observed (18) that acquisition of anoikis resistance by carcinoma cells is sufficient for their ability to grow as primary tumors. Thus, resistance of malignant cells to anoikis represents a major prerequisite for tumor progression (4;19;20). Hence, anoikis resistance of cancer cells may serve as a novel therapeutic target. However, molecular mechanisms that control anoikis in normal and cancer cells are understood at best in part.

Adherent cells are attached to the ECM via integrin receptors (21). Detachment-induced disengagement of integrins causes changes in the activity of various protein kinases, such as inhibition of c-Src (8) or activation of p38 mitogen-activated protein (MAP) kinase (22). These changes alter levels and/or activity of proteins that control cell survival, including proteins composing cellular apoptotic machinery.

One known apoptotic pathway involves the release of mitochondrial molecules such as cytochrome c, Smac and Omi/HtrA2 into the cytoplasm, which results in the activation cysteine proteases of the caspase family, such as the initiator caspase-9. Once activated, caspase-9 in turn triggers executioner caspases (23-28), which then cleave vital cellular targets and cause apoptosis (13;29). Caspases can be inhibited by the IAP family members, such as cIAP1, 2 and XIAP (30-32). Upon release from the mitochondria, Smac and Omi inactivate IAPs and trigger caspases (23-27;33;34). The release of the mitochondrial factors can be stimulated or blocked by pro-(Bak, Bax etc.) and anti-apoptotic (Bcl-2, Bcl-X_L etc.) proteins of the Bcl-2 family, respectively (35). Caspases can also be activated by another pathway that is induced by the activation of death receptors, such as Fas, by their ligands (Fas ligand) (36-39). Death receptors in turn activate initiator caspases 8 and 10, which then trigger the effector caspases (depending on the circumstances, either directly or by promoting the release of mitochondrial factors into the cytoplasm) and thus induce apoptosis.

We found so far that anoikis of intestinal epithelial cells is driven by detachment-induced downregulation of Bcl-X_L and subsequent release of Omi into the cytoplasm (28). We observed that, in addition, anoikis of these cells is mediated by detachment-dependent p38 MAP kinase-driven upregulation of Fas ligand (22).

Ras is a GTPase that is activated by receptor tyrosine kinases in response to diverse mitogenic signals (40). Activated Ras triggers multiple downstream pathways mediated by signaling molecules, such as Raf, Ral guanine nucleotide exchange factors (RalGEFs) and phosphoinositide 3-OH kinase (PI 3-kinase) (40). Some of these events promote changes in the expression of various genes. Ultimately, Ras-induced signaling mechanisms control proliferation, survival, and other critical cellular functions (41). Oncogenic mutations of *ras* often occur in numerous human cancers, including colorectal carcinoma (42;43).

Oncogenic *ras* is an efficient inhibitor of anoikis (28;44). According to our studies, Ras blocks anoikis of intestinal epithelial cells by triggering a network of anti-apoptotic signals, rather than by one mechanism. So far we have been able to identify some of the elements of this network. We have found that Ras blocks anoikis of intestinal epithelial cells by preventing detachment-induced downregulation of Bcl-X_L (12), by downregulating Bak (13) and by upregulating cIAP2 and XIAP (44). Importantly, we established that disruption of the effects of Ras on Bak and Bcl-X_L partially blocked anoikis resistance of *ras*-transformed cells *in vitro* and partly suppressed their tumorigenicity *in vivo* (12;13).

Whether or not all critical elements of the *ras*-activated network of anti-anoikis signals have been identified is not known. Furthermore, mechanisms linking Ras with those components of this network that have already been identified are understood poorly. Thus, which mediators (or their combination) of the anti-anoikis effect of Ras represent optimal targets for treatment aimed at the suppression of *ras*-driven anoikis resistance of cancer cells remains to be established.

In an effort to further understand the mechanisms allowing Ras to block anoikis we found that oncogenic Ras downregulates the initiator caspase-2 in intestinal epithelial cells.

Caspase-2 is known to mediate apoptosis triggered by diverse stimuli (45). The mechanisms by which this protease contributes to the execution of cellular apoptotic program are not well understood. It was proposed in this regard that caspase-2 can mediate apoptosis either downstream of the death receptors in complex with an adapter molecule RAIDD or as a part of a complex containing protein PIDD (45). However, according to the studies based on cells derived from PIDD and RAIDD knockout mice, activation of this caspase by various stimuli is not affected by complete loss of PIDD and RAIDD (46). Thus, the ability to form complexes with the indicated molecules does not seem to be essential for caspase-2-dependent apoptosis. One property of caspase-2 that many authors agree on is the ability of this caspase to induce cell death by facilitating (via poorly understood mechanisms) mitochondrial outer membrane permeabilization and thus stimulating the release of various mitochondrial factors into the cytoplasm (47-50).

Given that, to our knowledge, the effect of *ras*-induced downregulation of caspase-2 on anoikis resistance of malignant intestinal epithelial cells has never been studied we explored the role of this downregulation in the ability of *ras* oncogene-carrying cells to resist anoikis. We found that *ras*-dependent reduction of caspase-2 expression in these cells is required for their anoikis resistance and their ability to form tumors *in vivo*.

EXPERIMENTAL PROCEDURES

Cell culture- The generation of the IEC clones expressing activated H-*ras* (1) has been previously described. Expression of H-*ras* in MT-*ras* cells was induced by adding 100 μ M ZnCl₂ and 2 μ M CdCl₂ to cells. Clones of *ras*-3 cells expressing exogenous caspase-2 were generated using methods that we previously described (51). All IEC clones were cultured in α -minimum essential medium containing 5% fetal bovine serum, 10 μ g/ml insulin, and 0.5% glucose. The DLD-1, DKS-8 and DKO-3 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum. For suspension cultures cells were plated above a layer of 1% sea plaque-agarose polymerized in α -minimum

essential medium or Dulbecco's modified Eagle's medium.

Expression vectors- The expression vector pEGFP-N1 carrying Green Fluorescent Protein (GFP) fused to the C-terminus of caspase-2 was used for the transient transfection experiments (52). This vector was kindly provided by Dr. S. Kumar, Centre for Cancer Biology, Adelaide, Australia. pGL3b expression vector carrying firefly luciferase gene under the control of the fragment of caspase-2 gene containing caspase-2 promoter (spanning the DNA fragment located between positions -3970 and -2595 of the caspase-2 gene) was previously described (53). pRL expression vector carrying the *Renilla* luciferase was kindly provided by Dr. P. Lee, Dalhousie University, Halifax, NS Canada. For the generation of clones of *ras*-3 cells constitutively expressing caspase-2 GFP-tagged caspase-2 cDNA was placed into BamHI/NotI sites of the pcDNA4-TO vector (Invitrogen).

qPCR- Total RNA was isolated by use of the RNeasy Plus Mini Kit (Qiagen). RNA (3 μ g) was subsequently reverse transcribed by using RNA to cDNA EcoDry Kit (Clontech). The resulting cDNA was mixed with Brilliant SYBR Green qPCR Master Mix (Stratagene) and respective primers. qPCR and respective data analysis was performed as described (54) by use of MX3000P instrument under the following conditions. Samples were subjected to 10 min pre-denaturation at 95°C and then 40 cycles: 30 sec at 95°C, 1 min at 55°C and 30 sec at 72°C, each cycle. Samples were further incubated for 1 min at 95°C, 30 sec at 55°C and 30 sec at 95°C for the dissociation curve generation. The data were analyzed by MxPro QPCR software (Stratagene). Primers used to amplify respective rat cDNA were as follows: caspase-2 forward primer: TACTGCTCACAACCCTCTCT, reverse primer: TATAGGCCACGTAGTGT. 18S rRNA forward primer: AGTTCGAGTTAAAAGC, reverse primer: ACTCAGC AGAGCATCGAG.

Caspase-2 promoter activity assay - Iec-18 and *ras*3 cells were co-transfected with an expression vector coding for firefly luciferase under the control of the caspase-2 gene fragment containing caspase-2 promoter (1 μ g) and an expression vector coding for *Renilla* luciferase (0.25 μ g) for 24h as previously described (28). Cells were then lysed and respective lysates were

assayed for luciferase activity by use of Dual-Luciferase Report Assay kit (Promega). In the experiments when only ras-3 cells were transfected (in case of treatment with Mek1 inhibitor PD98059) the cells were transfected with 1 ug caspase-2 promoter-containing expression vector indicated above for 24h, after which cells were further cultured for additional 24h in the presence of 25uM PD98059 and assayed for luciferase activity as described above.

Transient transfection of ras-2 cells with caspase-2 expression vector-Transient transfection of ras-3 cells with a GFP-caspase-2 expression vector was performed as previously described (28).

Western blot analysis-Western blot analysis was performed as described elsewhere (44). The following antibodies were used in this study: anti-caspase-2 (Santa Cruz Biotechnology in Fig. 1, 2, 6 and Alexis in Fig. 3, 4), anti- α -tubulin (Upstate), anti-CDK4 (Santa Cruz Biotechnology), anti- β -actin (Sigma), anti-p38 MAP kinase (Santa Cruz Biotechnology), anti-GAPDH (Sigma), anti-cytochrome c (Cell Signaling), anti-HtrA2/Omi (R & D Systems). When lanes were removed from Western blot images, and separate parts of an image were joined together, a short vertical black line was used to indicate where the image was cut.

RNA interference-All transfections with siRNAs were performed by using Lipofectamine 2000 (Invitrogen) as previously described (44). The sequences of the sense strands of the RNAs used in this study were as follows. Control RNA (siCONTROL non-targeting siRNA #1, Dharmacon): UAGCGACUAAACACAUCAAUU; caspase-2 siRNA-1: GCACUUCACUGGAGAGAAAUU; caspase-2 siRNA-2: UCACAACCUCUCUGAUUUU. FADD siRNA-1: GGA AAAGACUGGCCCGUGA; FADD siRNA-2: GG GAUUCAACUGUGUCUUU. All RNAs were from Dharmacon.

Gene Expression Array-The expression of mRNAs coding for regulators of apoptosis was assayed by the rat-specific array carrying respective cDNAs (SuperArray) according to manufacturer's instructions. Signals on the array were detected by ECL. The intensity of each signal was quantified by densitometry as described (44).

Isolation of GFP-positive cells by flow cytometry-Cells were trypsinized, washed with and resuspended in a phosphate buffered saline buffer containing 25mM Hepes, 1% BSA and 1mM EDTA). FACSaria (BD Biosciences) instrument was used for the isolation of GFP-positive cells.

Analysis of apoptosis by flow cytometry-Apoptosis Detection kit from Chemicon was used in the assay. Cells were harvested, washed with PBS, and re-suspended in binding buffer provided by the manufacturer at a concentration of 10^6 cells/ml. 200 μ l of cell suspension was then mixed with 4 μ l of Annexin V conjugated to Allophycocyanin and 2 μ l of propidium iodide (20 μ g/ml), and the resulting mixture was incubated for 15 minutes at room temperature. FACSCalibur system (BD Biosciences) was used for the analysis. AnnexinV-positive propidium iodide-negative cells were considered apoptotic.

The following assays were performed as we previously described: measurement of the ability of cells to form colonies in monolayer after being cultured in suspension (55), soft agar growth assay (13), *in vivo* tumorigenicity assay (12).

Preparation of cytosolic fraction-Preparation of cytosolic fraction was performed as described by us and others (28;56).

Statistical analysis- 2-tailed Student's t-test was used for assessing statistical significance of data.

RESULTS

Oncogenic Ras blocks caspase-2 expression in intestinal epithelial cells. In an effort to understand the mechanisms by which oncogenic Ras blocks anoikis we compared levels of mRNAs coding for 97 apoptosis regulators in detached spontaneously immortalized non-malignant highly anoikis-susceptible intestinal epithelial cells IEC-18 and a previously published anoikis-resistant tumorigenic clone of these cells ras-3 (1;12) constitutively expressing oncogenic H-ras by using the array carrying respective cDNAs. One ras-induced change observed by us was the downregulation of the mRNA coding for the pro-apoptotic protein caspase-2 (Fig. 1A).

Caspase-2 is an initiator caspase that mediates the execution of apoptosis through

poorly understood mechanisms (45). This protease has been recently proposed to be able to act as a tumor suppressor in various contexts (45). Since the role of *ras*-dependent downregulation of caspase-2 in the control of anoikis of intestinal epithelial cells by oncogenic Ras, to our knowledge, has never been investigated, we decided to explore this role in the present study.

We confirmed by qPCR that caspase-2 mRNA levels are significantly lower in ras-3 cells than in IEC-18 cells (Fig. 1B). We further found that ras-3 cells transfected with an expression vector carrying luciferase gene under the control of a previously characterized (53) fragment of caspase-2 gene containing caspase-2 promoter (spanning the DNA fragment located between positions -3970 and -2595 of the caspase-2 gene) (53) displayed a noticeably lower luciferase activity than the parental IEC-18 cells transfected with the same vector (Fig. 1C). Collectively, the data presented above (Fig. 1A-C) indicate that Ras blocks transcription of the caspase-2 gene in the intestinal epithelial cells.

We further observed that caspase-2 expression is significantly lower at the protein level in ras-3 and ras-4 (another published anoikis-resistant tumorigenic clone of IEC-18 cells) cells (1;12) compared to the parental IEC-18 cells regardless of whether these cells were attached to or detached from the ECM (Fig. 1D). To confirm that the downregulation of caspase-2 represents a direct consequence of the presence of oncogenic *ras* in IEC-18 cells we utilized a published clone of IEC-18 cells MT-*ras* that harbors exogenous activated mutant of H-*ras* under the control of Zn²⁺- and Cd²⁺-inducible metallothionein promoter (57). We found that treatment of MT-*ras* cells with Zn²⁺ and Cd²⁺ results in a significant inhibition of caspase-2 expression (Fig. 1E, left). By contrast, treatment of the parental IEC-18 cells with the indicated metal ions did not cause any downregulation of caspase-2 (Fig. 1E, right).

To establish whether Ras can promote caspase-2 downregulation in human colon cancer cells, we utilized highly tumorigenic human colon carcinoma-derived cells DLD-1 carrying one allele of oncogenic K-*ras* and derivatives of these cells DKO-3 and DKS-8, in which the mutant K-*ras* allele had been disrupted by homologous recombination (58). We and others found previously that both oncogenic K-*ras*-deprived

variants of DLD-1 cells are significantly more anoikis-susceptible (12) and much less tumorigenic (58) than oncogenic *ras*-harboring DLD-1 cells. As shown in Fig. 1F, we observed that DLD-1 cells carry much lower amounts of caspase-2 than the mutant K-*ras*-knockout cells DKS-8 and DKO-3. Thus, *ras* oncogene downregulates caspase-2 in malignant intestinal epithelial cells.

Caspase-2 contributes to the execution of anoikis of intestinal epithelial cells. To test whether caspase-2 plays a role in the execution of anoikis of intestinal epithelial cells we ablated this caspase in IEC-18 cells by using two separate small interfering RNAs (siRNAs) targeted to different regions of caspase-2 mRNA (Fig. 2A, D). To assess the effect of enforced caspase-2 downregulation on the viability of detached cells we used a clonogenic cell survival assay that we often utilized in the past for measuring anoikis (28;55). In the course of the assay cells transfected with a control or caspase-2-specific siRNAs were cultured detached from the ECM, re-plated in monolayer, cells that remained viable after detachment were allowed to form colonies, and the resulting colonies were counted. We found that loss of caspase-2 significantly increases the viability of IEC-18 cells following detachment from the ECM (Fig. 2B, E). We further observed that enforced caspase-2 downregulation noticeably reduces the ability of detached IEC-18 cells to bind Annexin V (Fig. 2C, F) (this ability is one of the well established hallmarks of apoptosis (59)). Thus caspase-2 contributes to the execution of anoikis of non-malignant intestinal epithelial cells.

ras-induced downregulation of caspase-2 is required for anoikis resistance of ras-transformed intestinal epithelial cells. To address the role of *ras*-induced downregulation of caspase-2 in anoikis resistance of *ras*-transformed intestinal epithelial cells we decided to reverse this downregulation. To this end, we transiently transfected ras-3 (an oncogenic *ras*-expressing derivative of IEC-18 cells, see Fig. 1) with green fluorescent protein (GFP)-tagged caspase-2 (others established that caspase-2 remains active in the presence of the GFP tag (52)). We then isolated GFP-positive ras-3 cells by flow cytometry and confirmed that total amount of caspase-2 in the resulting cells is significantly higher than that in the cells transfected with a control vector carrying

GFP alone and does not exceed that in the parental IEC-18 cells (Fig. 3A). Thus, the subsequently observed effects of caspase-2 on apoptosis of ras-3 cells were not due to the presence of abnormally high amounts of this caspase in the indicated cells. We then tested the ability of GFP- and caspase-2-GFP-transfected cells to bind Annexin V and found that exogenous caspase-2 significantly increases the susceptibility of detached (Fig. 3C) but not that of the attached (Fig. 3B) ras-3 cells to apoptosis.

The ability of cells to grow in the absence of adhesion to the ECM as colonies in soft agar represents one of the most stringent criteria for malignant transformation that are presently being used (6;7). We thus tested whether the reversal of the effect of oncogenic Ras on caspase-2 blocks the long-term growth of the *ras*-transformed in soft agar. Since transient transfection of ras-3 cells with a caspase-2 expression vector was not optimal for these type of studies, we generated three clones of ras-3 cells, ras-casp2-2, ras-casp2-3 and ras-casp2-4 expressing ectopic caspase-2 in a constitutive manner. Again, we found that the total amount of caspase-2 in each of these clones was significantly higher than that in a vector control clone (ras-control) but did not exceed that in the parental IEC-18 cells (Fig. 4A). We further observed that exogenous caspase-2 had a relatively small effect on the ability of attached cells to form colonies (Fig. 4B) but noticeably blocked their clonogenicity in soft agar, when these cells were detached from the ECM (Fig. 4C). As expected, ectopic caspase-2, when expressed in ras-3 cells in a constitutive manner, also significantly increased their ability to bind Annexin V following detachment (Fig. 4D-F). Based on the data presented above, we concluded that *ras*-induced downregulation of caspase-2 is required for the ability of oncogenic Ras to protect intestinal epithelial cells from anoikis.

We found previously that one of the mechanisms of anoikis of intestinal epithelial cells, including IEC-18 cells, is mediated by Fas ligand, a pro-apoptotic protein that exerts its effect on cells via an adapter molecule FADD (22). We observed in the present study that ablation of FADD (Fig. 5A) in ras-casp2-4 cells (a clone of ras-3 cells expressing ectopic caspase-2, see Fig. 4) did not block their apoptosis following detachment (Fig. 5B, C). Thus, the pro-apoptotic

signaling pathway driven by Fas ligand and FADD does not appear to be required for the ability of caspase-2 to induce anoikis of ras-transformed intestinal epithelial cells.

ras-induced downregulation of caspase-2 prevents the release of mitochondrial apoptosis-inducing proteins into the cytoplasm of intestinal epithelial cells following detachment. Molecular events involved in caspase-2-initiated apoptosis are not well understood (45). It is however thought that one way by which this initiator caspase promotes cell death is via facilitating (through yet unknown mechanisms) the release of the pro-apoptotic mitochondrial factors into the cytoplasm, where they activate various elements of the cellular pro-apoptotic program (47-50). We found previously in this regard that detachment of intestinal epithelial cells (including IEC-18 cells) promotes the release of the mitochondrial proteins, such as cytochrome c and Omi/HtrA2 in the cytoplasm and that oncogenic Ras inhibits these events (28). Furthermore, we demonstrated that Ras-induced inhibition of detachment-induced release of the mitochondrial proteins, such as Omi, is required for the ability of oncogenic Ras to suppress anoikis (28). We thus tested whether the reversal of *ras*-induced downregulation of caspase-2 contributes to the ability of Ras to block the release of cytochrome c and Omi into the cytoplasm of detached cells. As shown in Figure 6A, B, detached cells ras-casp2-2, ras-casp2-3 and ras-casp2-4 expressing ectopic caspase-2 displayed significantly higher amounts of both cytochrome c and Omi in the cytoplasm than the respective control clone ras-control. In agreement with a notion that caspase-2 can mediate the release of the indicated pro-apoptotic factors in detached cells, we found that ablation of caspase-2 in the parental IEC-18 cells by RNAi (see Fig. 2) blocks such release (Fig. 6C, D). In summary, our data indicate that *ras*-induced downregulation of caspase-2 prevents the release of cytochrome c and Omi into the cytoplasm of intestinal epithelial cells following their detachment.

The activity of protein kinase Mek is required for ras-induced downregulation of caspase-2 in intestinal epithelial cells. Ras is known to be able to activate numerous signaling pathways, including those mediated by the sequential induction of protein kinases Raf, Mek and Erk (40). In an effort to identify the signaling

mechanism by which Ras downregulates caspase-2 in intestinal epithelial cells we found that treatment of ras-3 cells with PD98059, a specific and widely used small molecule inhibitor of Mek (60), resulted in a significant increase of the caspase-2 promoter activity (Fig. 7A), a noticeable upregulation of caspase-2 mRNA (Fig. 7B), and that of caspase-2 protein (Fig. 7C). By contrast, treatment with LY294002, an inhibitor of PI 3-kinase (61), another major mediator of Ras signaling (40), did not trigger caspase-2 protein upregulation in these cells (not shown). We reasoned that if the effect of Ras on caspase-2 is mediated by Mek and if this effect contributes to *ras*-induced anoikis resistance of intestinal epithelial cells, then inhibitors of Mek, such as PD98059 should promote anoikis of *ras*-transformed cells. Indeed, we found that treatment with PD98059 did not result in significant apoptosis of attached ras-3 cells but caused a noticeable increase of death of these cells when they were detached from the ECM (Fig. 7D). Collectively, these data are consistent with a scenario, according to which *ras*-induced downregulation of caspase-2 is mediated by Mek.

ras-induced downregulation of caspase-2 is required for the ability of *ras*-transformed intestinal epithelial cells to form tumors *in vivo*. Normal intestinal epithelium exists *in vivo* as a single layer, whereas primary tumors as well as tumors formed by cancer cells subcutaneously injected in mice, a model that is often used for studying tumorigenesis, tend to grow as three-dimensional masses. The results of several studies, including ours, indicate that anoikis resistance of cancer cells is required for the ability of these cells to form tumors following subcutaneous injection in mice (11-15). Given that *ras*-induced downregulation of caspase-2 is required for the ability *ras*-transformed cells to resist anoikis and grow without being attached to the ECM, we decided to test whether this downregulation contributes to the ability of the indicated cells to form tumors in mice. As shown in Figure 8, the sizes of tumors formed by cells *ras*-casp2-2, *ras*-casp2-3 and *ras*-casp2-4 expressing ectopic caspase-2 were significantly smaller at all times of the assay than those of the tumors formed by respective control cells. Thus, *ras*-induced downregulation of caspase-2 does contribute to the *in vivo* tumorigenicity of these cells.

In summary, we have identified a novel mechanism by which oncogenic Ras blocks anoikis of intestinal epithelial cells, allows them to grow in an anchorage-independent manner within three-dimensional multicellular masses and enables them to form tumors. This mechanism is driven by *ras*-induced downregulation of caspase-2.

DISCUSSION

We have identified in this study a novel mechanism by which oncogenic Ras promotes anoikis resistance of intestinal epithelial cells. This mechanism involves *ras*-dependent downregulation of caspase-2. We found previously that anoikis of non-malignant intestinal epithelial cells is driven by detachment-induced downregulation of Bcl-X_L (12) and subsequent release of the mitochondrial factors, such as HtrA2/Omi, into the cytoplasm (28). We have demonstrated in this study that in addition, anoikis of these cells is mediated (via mechanisms that remain to be established) by caspase-2, a protease that according to this (Fig. 6) and previous (47-50) studies does have the ability to increase the permeability of the mitochondria to the pro-apoptotic factors. The present as well as our previous studies indicate that, oncogenic Ras has the ability to block this network of pro-anoikis signals in detached cells by activating a network of the anti-anoikis signals. We found in the past that two important elements of this network are the mechanisms involving *ras*-induced downregulation of Bak (13) and inhibition of detachment-induced downregulation of Bcl-X_L (12). We show here that one additional, mechanism by which Ras prevents the release of cytotoxic factors, such as cytochrome c and Omi from the mitochondria is driven by *ras*-dependent downregulation of caspase-2.

Our data suggest that Ras downregulates caspase-2 in intestinal epithelial cells by triggering a protein kinase Mek, an inducer of the mitogen activated protein kinases and a mediator of Ras signaling whose activity is known to be stimulated by a Ras binding partner protein kinase Raf (40). To our knowledge, the present study, for the first time demonstrates that the indicated signaling pathway can block anoikis downstream of Ras in intestinal epithelial cells. Of note, we observed in

the past that *ras*-induced downregulation of Bak, the second mechanism by which oncogenic Ras blocks the release of the pro-apoptotic mitochondrial factors in these cells and their subsequent anoikis, is driven by PI 3-kinase (13) which represents another important mediator of Ras signaling (40). Therefore, it is the activation of both major Ras-induced signaling pathways, one mediated by Mek and another one, controlled by PI 3-kinase, that contribute to the indicated effects of Ras in intestinal epithelial cells.

Interestingly, we have found that exogenous caspase-2 promoted a much more noticeable apoptosis of the *ras*-transformed cells when they were detached from the ECM than in the attached cells (see Figures 3 and 4). These data suggest that a threshold caspase-2 concentration is required for the induction of apoptosis by signals that are induced by loss of adhesion of intestinal epithelial cells to the ECM. However, it seems likely that when caspase-2 is downregulated in these cells (e.g. in response to the expression of oncogenic *ras*), signals that are induced by detachment of the indicated cells become by themselves insufficient for stimulating apoptosis.

We found in this study that *ras*-induced downregulation of caspase-2 is required for the ability of *ras*-transformed intestinal epithelial cells to form tumors *in vivo*. These findings well agree with what is known about the role of anoikis resistance of cancer in growth of tumors formed by malignant cells. It has been well established in this regard that normal intestinal epithelium exists *in vivo* as a single layer, whereas primary human tumors as well as tumors formed by cancer cells that have been subcutaneously injected into mice, a model that we have used in this study, typically form three-dimensional masses. Perhaps not by coincidence, the ability of cancer cells to grow in a three-dimensional anchorage-independent manner in soft agar has served as a “gold standard” for malignant transformation for several decades (6), and cells that are capable of this growth can usually form subcutaneous tumors (7;12). Moreover, treatment that blocks resistance of cancer cells to detachment-induced death is known to inhibit their ability to form such tumors (11-14). We found in the past, for example, that the reversal of *ras*-induced downregulation of Bak

(13) or the ablation of Bcl-X_L (12) in the *ras*-transformed intestinal epithelial cells enhances their susceptibility to detachment-induced death and blocks their ability to form subcutaneous tumors in mice. In addition, we found that variants of the poorly tumorigenic intestinal epithelial cells selected for increased anoikis resistance acquire the capacity for forming such tumors (18). The results of this study indicate that the reversal of *ras*-induced downregulation of caspase-2 in malignant intestinal epithelial cells represents a relatively efficient approach for blocking growth of tumors formed by these cells.

The fact that caspase-2 can suppress anoikis and three-dimensional tumor growth is consistent with several other studies pointing at the tumor suppression function for this caspase. It was shown in this regard that mouse embryonic fibroblasts (MEFs) derived from caspase-2 knockout mice, when transformed with oncogenes, grow faster in monolayer culture as well as in soft agar and are more tumorigenic in mice than similarly transformed wild type MEFs (62). In this case, however, loss of caspase-2 seemed to accelerate proliferation of transformed MEFs, rather than block their anoikis (caspase-2 is known to have the ability to block the cell cycle progression under certain circumstances (45)). The fact that caspase-2 mediates proliferation, rather than anoikis, of MEFs is not surprising, in view of the fact that MEFs tend not to be prone to anoikis unless they are deprived of growth factors (63;64). It is also known that caspase-2 expression is frequently reduced in human gastric tumors when compared to normal gastric mucosa (65). Furthermore, caspase-2 was found to be significantly underexpressed in metastatic brain tumors (66). Finally, caspase-2-deficient mice were demonstrated to be noticeably more susceptible to Myc-induced lymphoma than the respective wild type mice (62).

In summary, our data indicate that the anti-apoptotic mechanism triggered by *ras*-induced downregulation of caspase-2 represents an important novel element of the signaling network via which oncogenic Ras blocks anoikis and promotes three-dimensional growth of tumors formed by malignant intestinal epithelial cells.

Reference List

1. Rak, J., Mitsuhashi, Y., Erdos, V., Huang, S. N., Filmus, J., and Kerbel, R. S. (1995) *J.Cell Biol.* **131**, 1587-1598
2. Frisch, S. M. and Francis, H. (1994) *J.Cell Biol.* **124**, 619-626
3. Ljubimov, A. V., Bartek, J., Couchman, J. R., Kapuller, L. L., Veselov, V. V., Kovarik, J., Perevoshchikov, A. G., and Krutovskikh, V. A. (1992) *Int.J.Cancer* **50**, 562-566
4. Douma, S., Van Laar, T., Zevenhoven, J., Meuwissen, R., Van Garderen, E., and Peeper, D. S. (2004) *Nature* **430**, 1034-1039
5. Berezovskaya, O., Schimmer, A. D., Glinskii, A. B., Pinilla, C., Hoffman, R. M., Reed, J. C., and Glinsky, G. V. (2005) *Cancer Res.* **65**, 2378-2386
6. Freedman, V. H. and Shin, S. I. (1974) *Cell* **3**, 355-359
7. Lim, K. H., Baines, A. T., Fiordalisi, J. J., Shipitsin, M., Feig, L. A., Cox, A. D., Der, C. J., and Counter, C. M. (2005) *Cancer Cell* **7**, 533-545
8. Rosen, K., Coll, M. L., Li, A., and Filmus, J. (2001) *J.Biol.Chem.* **276**, 37273-37279
9. Li, H., Ray, G., Yoo, B. H., Erdogan, M., and Rosen, K. V. (2009) *J.Biol.Chem.* **284**, 2012-2022
10. Lu, Y., Lin, Y. Z., LaPushin, R., Cuevas, B., Fang, X., Yu, S. X., Davies, M. A., Khan, H., Furui, T., Mao, M., Zinner, R., Hung, M. C., Steck, P., Siminovitch, K., and Mills, G. B. (1999) *Oncogene* **18**, 7034-7045
11. Duxbury, M. S., Ito, H., Zinner, M. J., Ashley, S. W., and Whang, E. E. (2004) *Oncogene* **23**, 1448-1456
12. Rosen, K., Rak, J., Leung, T., Dean, N. M., Kerbel, R. S., and Filmus, J. (2000) *J.Cell Biol.* **149**, 447-456
13. Rosen, K., Rak, J., Jin, J., Kerbel, R. S., Newman, M. J., and Filmus, J. (1998) *Curr.Biol.* **8**, 1331-1334
14. Scotlandi, K., Maini, C., Manara, M. C., Benini, S., Serra, M., Cerisano, V., Strammiello, R., Baldini, N., Lollini, P. L., Nanni, P., Nicoletti, G., and Picci, P. (2002) *Cancer Gene Ther.* **9**, 296-307
15. Frankel, A., Rosen, K., Filmus, J., and Kerbel, R. S. (2001) *Cancer Res.* **61**, 4837-4841
16. Duxbury, M. S., Ito, H., Zinner, M. J., Ashley, S. W., and Whang, E. E. (2004) *Oncogene* **23**, 465-473
17. Jiang, K., Sun, J., Cheng, J., Djeu, J. Y., Wei, S., and Sebt, S. (2004) *Mol.Cell Biol.* **24**, 5565-5576
18. Derouet, M., Wu, X., May, L., Hoon, Y. B., Sasazuki, T., Shirasawa, S., Rak, J., and Rosen, K. V. (2007) *Neoplasia.* **9**, 536-545

19. Evan, G. I. and Vousden, K. H. (2001) *Nature* **411**, 342-348
20. Jacks, T. and Weinberg, R. A. (2002) *Cell* **111**, 923-925
21. Ruoslahti, E. and Reed, J. C. (1994) *Cell* **77**, 477-478
22. Rosen, K., Shi, W., Calabretta, B., and Filmus, J. (2002) *J.Biol.Chem.* **277**, 46123-46130
23. Li, P., Nijhawan, D., Budihardjo, I., Srinivasula, S. M., Ahmad, M., Alnemri, E. S., and Wang, X. (1997) *Cell* **91**, 479-489
24. Du, C., Fang, M., Li, Y., Li, L., and Wang, X. (2000) *Cell* **102**, 33-42
25. Verhagen, A. M., Ekert, P. G., Pakusch, M., Silke, J., Connolly, L. M., Reid, G. E., Moritz, R. L., Simpson, R. J., and Vaux, D. L. (2000) *Cell* **102**, 43-53
26. Suzuki, Y., Imai, Y., Nakayama, H., Takahashi, K., Takio, K., and Takahashi, R. (2001) *Mol.Cell* **8**, 613-621
27. Verhagen, A. M., Silke, J., Ekert, P. G., Pakusch, M., Kaufmann, H., Connolly, L. M., Day, C. L., Tikoo, A., Burke, R., Wrobel, C., Moritz, R. L., Simpson, R. J., and Vaux, D. L. (2002) *J.Biol.Chem.* **277**, 445-454
28. Liu, Z., Li, H., Derouet, M., Berezkin, A., Sasazuki, T., Shirasawa, S., and Rosen, K. (2006) *J.Biol.Chem.* **281**, 14738-14747
29. Salvesen, G. S. and Dixit, V. M. (1997) *Cell* **91**, 443-446
30. Li, F. (2003) *J.Cell Physiol* **197**, 8-29
31. Deveraux, Q. L. and Reed, J. C. (1999) *Genes Dev.* **13**, 239-252
32. LaCasse, E. C., Baird, S., Korneluk, R. G., and MacKenzie, A. E. (1998) *Oncogene* **17**, 3247-3259
33. Gottfried, Y., Rotem, A., Lotan, R., Steller, H., and Larisch, S. (2004) *EMBO J.* **23**, 1627-1635
34. Olson, M. and Kornbluth, S. (2001) *Curr.Mol.Med.* **1**, 91-122
35. Tsujimoto, Y. (2003) *J.Cell Physiol* **195**, 158-167
36. Locksley, R. M., Killeen, N., and Lenardo, M. J. (2001) *Cell* **104**, 487-501
37. Ashkenazi, A. (2002) *Nat.Rev.Cancer* **2**, 420-430
38. Algeciras-Schimmich, A., Shen, L., Barnhart, B. C., Murmann, A. E., Burkhardt, J. K., and Peter, M. E. (2002) *Mol.Cell Biol.* **22**, 207-220
39. Medema, J. P., Scaffidi, C., Kischkel, F. C., Shevchenko, A., Mann, M., Krammer, P. H., and Peter, M. E. (1997) *EMBO J.* **16**, 2794-2804

40. Shields, J. M., Pruitt, K., McFall, A., Shaub, A., and Der, C. J. (2000) *Trends Cell Biol.* **10**, 147-154
41. Khosravi-Far, R., Campbell, S., Rossman, K. L., and Der, C. J. (1998) *Adv.Cancer Res.* **72**, 57-107
42. Bos, J. L., Fearon, E. R., Hamilton, S. R., Verlaan-de Vries, M., van Boom, J. H., van der Eb, A. J., and Vogelstein, B. (1987) *Nature* **327**, 293-297
43. Barbacid, M. (1987) *Annu.Rev.Biochem.* **56**, 779-827
44. Liu, Z., Li, H., Derouet, M., Filmus, J., LaCasse, E. C., Korneluk, R. G., Kerbel, R. S., and Rosen, K. V. (2005) *J.Biol.Chem.* **280**, 37383-37392
45. Kumar, S. (2009) *Nat.Rev.Cancer* **9**, 897-903
46. Manzl, C., Krumschnabel, G., Bock, F., Sohm, B., Labi, V., Baumgartner, F., Logette, E., Tschopp, J., and Villunger, A. (2009) *J.Cell Biol.* **185**, 291-303
47. Guo, Y., Srinivasula, S. M., Druilhe, A., Fernandes-Alnemri, T., and Alnemri, E. S. (2002) *J.Biol.Chem.* **277**, 13430-13437
48. Lassus, P., Opitz-Araya, X., and Lazebnik, Y. (2002) *Science* **297**, 1352-1354
49. Robertson, J. D., Enoksson, M., Suomela, M., Zhivotovsky, B., and Orrenius, S. (2002) *J.Biol.Chem.* **277**, 29803-29809
50. Wagner, K. W., Engels, I. H., and Deveraux, Q. L. (2004) *J.Biol.Chem.* **279**, 35047-35052
51. Yoo, B. H., Wu, X., Li, Y., Haniff, M., Sasazuki, T., Shirasawa, S., Eskelinen, E. L., and Rosen, K. V. (2010) *J.Biol.Chem.* **285**, 5438-5449
52. Colussi, P. A., Harvey, N. L., and Kumar, S. (1998) *J.Biol.Chem.* **273**, 24535-24542
53. Logette, E., Wotawa, A., Solier, S., Desoche, L., Solary, E., and Corcos, L. (2003) *Oncogene* **22**, 935-946
54. Schmittgen, T. D. and Livak, K. J. (2008) *Nat.Protoc.* **3**, 1101-1108
55. Liu, Z., Li, H., Wu, X., Yoo, B. H., Yan, S. R., Stadnyk, A. W., Sasazuki, T., Shirasawa, S., LaCasse, E. C., Korneluk, R. G., and Rosen, K. V. (2006) *Oncogene* **25**, 7680-7690
56. Rytomaa, M., Lehmann, K., and Downward, J. (2000) *Oncogene* **19**, 4461-4468
57. Filmus, J., Robles, A. I., Shi, W., Wong, M. J., Colombo, L. L., and Conti, C. J. (1994) *Oncogene* **9**, 3627-3633
58. Shirasawa, S., Furuse, M., Yokoyama, N., and Sasazuki, T. (1993) *Science* **260**, 85-88
59. van Engeland, M., Nieland, L. J., Ramaekers, F. C., Schutte, B., and Reutelingsperger, C. P. (1998) *Cytometry* **31**, 1-9

60. Alessi, D. R., Cuenda, A., Cohen, P., Dudley, D. T., and Saltiel, A. R. (1995) *J.Biol.Chem.* **270**, 27489-27494
61. Vlahos, C. J., Matter, W. F., Hui, K. Y., and Brown, R. F. (1994) *J.Biol.Chem.* **269**, 5241-5248
62. Ho, L. H., Taylor, R., Dorstyn, L., Cakouros, D., Bouillet, P., and Kumar, S. (2009) *Proc.Natl.Acad.Sci.U.S A* **106**, 5336-5341
63. Meredith, J. E. and Schwartz, M. A. (1997) *Trends in Cell Biology* **7**, 146-150
64. Zouq, N. K., Keeble, J. A., Lindsay, J., Valentijn, A. J., Zhang, L., Mills, D., Turner, C. E., Streuli, C. H., and Gilmore, A. P. (2009) *J.Cell Sci.* **122**, 357-367
65. Yoo, N. J., Lee, J. W., Kim, Y. J., Soung, Y. H., Kim, S. Y., Nam, S. W., Park, W. S., Lee, J. Y., and Lee, S. H. (2004) *APMIS* **112**, 330-335
66. Zohrabian, V. M., Nandu, H., Gulati, N., Khitrov, G., Zhao, C., Mohan, A., Demattia, J., Braun, A., Das, K., Murali, R., and Jhanwar-Uniyal, M. (2007) *Oncol.Rep.* **18**, 321-328

FOOTNOTES

This study was supported by a grant from Canadian Institutes of Health Research. We are grateful to Drs. S. Kumar and P. Lee for the materials provided to us and to Mrs. O. Vitviskaia for technical assistance.

FIGURE 1 Oncogenic *ras* downregulates caspase-2 in intestinal epithelial cells. *A*, IEC-18 and a H-*ras*-transformed clone of these cells *ras*-3 were cultured detached from the ECM for 24h and the expression of caspase-2 mRNA (along with other regulators of apoptosis) was analyzed in these cells by use of the rat-specific array carrying respective cDNAs (SuperArray). Signals on the array were detected by ECL. The intensity of each signal was quantified by densitometry and normalized by the levels of Ube2i mRNA that served as a loading control. *B*, Indicated cell lines were analyzed for caspase-2 mRNA expression by qPCR. The observed caspase-2 mRNA levels were normalized by the levels of 18S rRNA which were also determined by qPCR. The resulting levels of caspase-2 mRNA in IEC-18 cells were arbitrarily designated as 1.0. The data represent the average of three independent experiments plus the SD. *C*, Indicated cell lines were transiently transfected with expression vector coding for firefly luciferase under the control of the caspase-2 gene fragment containing caspase-2 promoter and an expression vector coding for *Renilla* luciferase. The intensity of respective signals corresponding to firefly luciferase activity was normalized by those of *Renilla* luciferase. The resulting numbers obtained for IEC-18 cells were arbitrarily designated as 1.0. The data represent the average of three independent experiments plus the SD. The values observed for IEC-18 cells and marked with an asterisk in (*A-C*) were significantly ($p < 0.05$) different from those observed for *ras*-3 cells. *D*, IEC-18 cells and two independently derived H-*ras*-transformed clones of these cells *ras*-3 and *ras*-4 were cultured attached (att) to and detached (det) from the ECM for 24h and assayed for the expression of caspase-2 by Western blot. *E*, Attached MT-*ras* cells (left) or IEC-18 cells (right) were cultured in the absence (-) and in the presence (+) of 100 μM Zn^{2+} and 2 μM Cd^{2+} for 24h (left) and assayed for the expression of caspase-2 by Western blot. *F*, Attached human colorectal carcinoma cells DLD-1 and their K-*ras* knock-out derivatives DKO-3 and DKS-8 were assayed for the expression of caspase-2 by Western blot. The membranes in *D* and *E* were re-probed with a CDK-4 antibody and the membranes in *F* with an anti-p38 MAP kinase as loading controls.

FIGURE 2 Caspase-2 is required for anoikis of intestinal epithelial cells. *A*, *D*, IEC-18 cells were transfected with a control RNA (contRNA) or caspase-2-specific siRNA1 (casp2siRNA-1, *A*) or caspase-2-specific siRNA2 (casp2siRNA-2, *D*) and assayed for caspase-2 expression by Western blot. CDK-4 was used as a loading control. *B*, *E*, Cells were subsequently placed in suspension for 24 h and then re-plated in monolayer. The number of viable cells after detachment was calculated as a number of colonies formed seven days later by the cells that survived after being cultured in suspension. The data represent the average of the triplicates plus the SD. This experiment was repeated twice with similar results. *C*, *F*, Cells transfected as in (*A*, *D*, respectively) were placed in suspension for 24h and assayed for Annexin V binding by flow cytometry. The data represent the average of three experiments plus the S.E. The values marked with an asterisk were significantly ($p < 0.05$) different than those derived from the respective control experiments.

FIGURE 3 *ras*-induced downregulation of caspase-2 is required for anoikis resistance of malignant intestinal epithelial cells. *A*, *ras*-3 cells were transfected either with a control vector carrying GFP alone or with an expression vector carrying GFP-tagged caspase-2 and assayed for caspase-2 expression along with the parental IEC-18 cells by Western blot. A caspase-2-specific antibody was used in the assay. The positions of endogenous (endog.) caspase-2 and exogenous GFP-tagged caspase-2 (caspase-2-GFP) on the gel are indicated. CDK-4 was used as a loading control. *B*, *C*, Cells processed as in (*A*) were cultured attached to (*B*) detached from (*C*) the ECM for 48h and analyzed for Annexin V binding by flow cytometry. The data represent the average of three (*B*) and two (*C*) independent experiments plus the SD. These experiments were repeated twice with similar results. The value marked with an asterisk was significantly ($p < 0.05$) higher than that derived from the respective control experiments.

FIGURE 4 *ras*-induced downregulation of caspase-2 is required for the ability of malignant intestinal epithelial cells to grow in an anchorage-independent manner. *A*, IEC-18 cells, clones of *ras*-3 cells *ras*-casp2-2, *ras*-casp2-3 and *ras*-casp2-4 (generated by transfection of *ras*-3 cells with a caspase-2-

GFP expression vector) expressing ectopic caspase-2 in a constitutive manner and a control clone of ras-3 cells (generated by transfection of ras-3 cells with a control vector) were assayed for caspase-2 expression by Western blot. A caspase-2-specific antibody was used in the assay. The positions of endogenous (endog.) caspase-2 and exogenous GFP-tagged caspase-2 (caspase-2-GFP) on the gel are indicated. CDK-4 was used as a loading control. *B, C*, Indicated cell lines were plated in monolayer (*B*) or in soft agar (*C*), and colonies formed by these cells were counted 7-10 days later. The number of colonies formed by the ras-control cells was arbitrarily designated as 1.0. The data represent the average of three independent experiments plus the SD. *D-F*, Indicated cell lines were assayed for Annexin V binding by flow cytometry. The data in (*D*) represent the average of two independent experiments plus the SD. The data in (*E, F*) represent the average of three independent experiments plus the SEM. Values marked with an asterisk were significantly ($p < 0.05$) different from those derived from the respective control experiments.

FIGURE 5 Apoptosis of detached *ras*-transformed intestinal epithelial cells expressing exogenous caspase-2 can not be blocked by the ablation of FADD. ras-casp2-4 cells (a clone of ras-3 cells expressing exogenous caspase-2) were transfected with a control RNA (contRNA) or FADD-specific siRNA1 (FADDsiRNA-1) or FADD-specific siRNA2 (FADDsiRNA-2) and assayed for FADD expression by Western blot. CDK-4 was used as a loading control. Cells transfected as in (*A*) were placed in suspension for 48h (*B*) or 72h (*C*) and assayed for Annexin V binding by flow cytometry. The data represent the average of two independent experiments plus the S.D.

FIGURE 6 *ras*-induced downregulation of caspase-2 prevents the release of cytochrome c and Omi/HtrA2 into the cytoplasm following detachment of intestinal epithelia cells. *A, B*, Indicated cell lines were cultured in suspension for 24h, cytosolic material was isolated from these cells and assayed for the presence of cytochrome c (*A*) and Omi (*B*) by Western blot. *C, D*, IEC-18 cells were transfected as in Fig. 2 with a control RNA (contRNA) or caspase-2-specific siRNA1 (casp2siRNA-1) or caspase-2-specific siRNA2 (casp2siRNA-2), cultured in suspension for 2h, cytosolic material was isolated from these cells and assayed for the presence of cytochrome c (*C*) and Omi (*D*) by Western blot. α -tubulin was used as a loading control.


FIGURE 7 The activity of protein kinase Mek is required for *ras*-induced downregulation of caspase-2 in intestinal epithelial cells. *A*, Cells were transiently transfected with expression vector coding for firefly luciferase under the control of the caspase-2 gene fragment containing caspase-2 promoter, treated with either DMSO (-) or 25 μ M PD98059 (+) for 24h and assayed for luciferase activity. The resulting numbers obtained for the untreated cells were arbitrarily designated as 1.0. The data represent the average of three independent experiments plus the SD. *B*, ras-3 cells were treated with PD98059 as in *A* and caspase-2 mRNA levels were measured in these cells by qPCR. The observed caspase-2mRNA levels were normalized by the levels of 18S rRNA which were also determined by qPCR. The resulting levels of caspase-2 mRNA DMSO-treated cells were arbitrarily designated as 1.0. The data represent the average of three independent experiments plus the SD. *C*, Cells treated with PD98059 as in *A* were assayed for caspase-2 expression by Western blot. GAPDH was used as a loading control. *D*, Cells were cultured in monolayer or suspension for 24 in the presence of either DMSO (-) or 25 μ M PD98059 (+) and analyzed for Annexin V binding by flow cytometry. The data represent the average of two independent experiments plus the SD. The values marked with an asterisk was significantly ($p < 0.05$) different from those derived from respective control experiments.

FIGURE 8 *ras*-induced downregulation of caspase-2 is required for the ability of malignant intestinal epithelial cells to form tumors in mice. The indicated cell lines were injected subcutaneously into nude mice, and tumor volumes were measured at the indicated time points. Four mice were injected with each cell line. Error bars represent the SEM. This experiment was repeated twice with similar


results. Asterisks indicate that values derived from the control experiments with ras-control cells were significantly ($p < 0.05$) higher than those observed for respective experiments with ras-casp2-2, ras-casp2-3 and ras-casp2-4 cells.

Figure 1


A.




B.




C.



D.



E.



F.






Figure 2


A.




B.




C.



D.



E.



F.






Figure 3

A.



B.



C.




Figure 4






Figure 5

A.



B.



C.




Figure 6









Figure 7


A.



B.



C.



D.




Figure 8

