

Effect of Nitrate and Acetylene on *nirS*, *cnorB*, and *nosZ* Expression and Denitrification Activity in *Pseudomonas mandelii*[∇]

Saleema Saleh-Lakha,^{1,2} Kelly E. Shannon,^{1,2} Sherri L. Henderson,¹ Bernie J. Zebarth,¹
David L. Burton,³ Claudia Goyer,^{1*} and Jack T. Trevors^{2*}

Potato Research Centre, Agriculture and Agri-Food Canada, Fredericton, New Brunswick E3B 4Z7, Canada¹; School of Environmental Sciences, University of Guelph, Guelph, Ontario N1G 2W1, Canada²; and Department of Environmental Sciences, Nova Scotia Agricultural College, Truro, Nova Scotia B2N 5E3, Canada³

Received 6 April 2009/Accepted 3 June 2009

Nitrate acts as an electron acceptor in the denitrification process. The effect of nitrate in the range of 0 to 1,000 mg/liter on *Pseudomonas mandelii* *nirS*, *cnorB*, and *nosZ* gene expression was studied, using quantitative reverse transcription-quantitative PCR. Denitrification activity was measured by using the acetylene blockage method and gas chromatography. The effect of acetylene on gene expression was assessed by comparing denitrification gene expression in *P. mandelii* culture grown in the presence or absence of acetylene. The higher the amount of NO_3^- present, the greater the induction and the longer the denitrification genes remained expressed. *nirS* gene expression reached a maximum at 2, 4, 4, and 6 h in cultures grown in the presence of 0, 10, 100, and 1,000 mg of KNO_3 /liter, respectively, while induction of *nirS* gene ranged from 12- to 225-fold compared to time zero. *cnorB* gene expression also followed a similar trend. *nosZ* gene expression did not respond to NO_3^- treatment under the conditions tested. Acetylene decreased *nosZ* gene expression but did not affect *nirS* or *cnorB* gene expression. These results showed that *nirS* and *cnorB* responded to nitrate concentrations; however, significant denitrification activity was only observed in culture with 1,000 mg of KNO_3 /liter, indicating that there was no relationship between gene expression and denitrification activity under the conditions tested.

Denitrification is a critical biological reaction in the global nitrogen cycle that involves the reduction of nitrate (NO_3^-) or nitrite (NO_2^-) to gaseous nitric oxide (NO), nitrous oxide (N_2O), or molecular nitrogen (N_2) under oxygen-limited conditions (26). The main environmental factors that control denitrification include the availability of substrates (i.e., NO_3^- or NO_2^-), oxygen limitation, carbon availability, pH, temperature, and the presence of denitrifying microorganisms (21). The presence of N-oxides, as electron acceptors, is essential in maintaining denitrification activity (9, 21, 24). Nitrate is an excellent choice as a parameter to study the effects on denitrification gene expression and activity, as opposed to NO_2^- , due to its prevalence in nature and lack of toxicity at high concentrations.

The effect of NO_3^- on denitrification in a pure culture has been studied through biochemical measurements of changes in NO_3^- , NO_2^- , and denitrification products (N_2O) (4, 5, 19, 22). Thomas et al. (22) measured denitrification in nonproliferating cell suspensions of *Pseudomonas aeruginosa* and *Pseudomonas fluorescens* and concluded that the optimum NO_3^- concentration for denitrification activity in both cultures was 20 mM. A separate study (5) investigated the rate of denitrification in *Pseudomonas stutzeri*, *P. aeruginosa*, and *Paracoccus denitrifi-*

cans and concluded that all three organisms had different denitrification rates and accumulated different amounts of NO_2^- , N_2O , and N_2 . Most research on denitrification gene expression in *Pseudomonas* spp. have focused on gene regulation (1, 9, 10); however, there are some studies that have investigated the effect of NO_3^- on denitrification gene expression. Denitrification in pure culture has been studied to understand the effect of NO_3^- (10, 14, 19) on denitrification gene expression. Körner and Zumft (14) examined the effect of various N-oxides on the production of denitrification enzymes in pure cultures of *P. stutzeri* and concluded that NO_3^- resulted in maximum synthesis of all denitrification reductases, compared to NO_2^- and N_2O . An investigation on the effect of NO_3^- on denitrification gene expression and activity in *P. mandelii*, a dominant culturable denitrifier isolated from an agricultural field in Canada (7), was limited to measuring *cnorB* gene expression in the presence (1,000 mg/liter) or absence of KNO_3 addition (19). Nitrate-amended cells expressed *cnorB* at a high level (average of 2.06×10^8 transcripts/ μg RNA) for 6 h and had significant N_2O accumulations in the presence of acetylene (101 μmol). When NO_3^- was not present, transient *cnorB* gene expression was observed for 3 h, beyond which transcript numbers declined to an average of 3.63×10^6 transcripts/ μg RNA, from 4 to 6 h. Moreover, insignificant denitrification was measured in cells grown in the absence of nitrate. Although the present study was useful in understanding the effect of NO_3^- on *cnorB* gene expression in *P. mandelii* culture with or without NO_3^- addition, the effect of different NO_3^- concentrations, including intermediate NO_3^- concentrations on *cnorB*, *nirS*, and *nosZ* denitrification genes, was not addressed. Many denitrification studies use the acetylene inhibition method to measure short-

* Corresponding author. Mailing address for C. Goyer: Potato Research Centre, Agriculture and Agri-Food Canada, Fredericton, New Brunswick E3B 4Z7, Canada. Phone: (506) 452-4851. Fax: (506) 452-3316. E-mail: claudia.goyer@agr.gc.ca. Mailing address for J. T. Trevors: School of Environmental Sciences, University of Guelph, Guelph, Ontario N1G 2W1, Canada. Phone: (519) 824-4120, ext. 53367. Fax: (519) 837-0442. E-mail: jtrevors@uoguelph.ca.

[∇] Published ahead of print on 12 June 2009.

term denitrification activity in soil (6, 11, 15, 17, 18, 20, 23) and pure cultures (3, 19, 25). Acetylene blocks the action of nitrous oxide reductase enzyme, thereby inhibiting the reduction of N_2O to N_2 (3, 12, 25). As a result, N_2O becomes the terminal product of denitrification and can be quantified by gas chromatography. Although the mechanism of action of acetylene has been widely investigated and established, the effect of acetylene on denitrification gene expression has not yet been determined.

The objective of this research was to determine the effect of various NO_3^- concentrations (0, 10, 100, and 1,000 mg/liter) on growth, denitrification gene expression (*nirS*, *cnorB*, and *nosZ*), and denitrification activity in a pure culture of *P. mandelii*. The effect of acetylene on denitrification gene expression (*nirS*, *cnorB*, and *nosZ*) in *P. mandelii* cultures was also assessed. It was hypothesized that (i) increasing concentrations of NO_3^- would result in improved growth, in a longer period of time during which denitrification genes are highly expressed and, subsequently, in higher denitrification activity as measured by N_2O product in the presence of acetylene and (ii) the presence of acetylene would not influence denitrification gene expression.

MATERIALS AND METHODS

Bacterial strains and growth conditions. *P. mandelii* strain PD30 was cultured in tryptic soy broth (TSB) medium (Difco, Bacto, Dickinson and Company, Sparks, MA) at 30°C and maintained on tryptic soy agar plates as previously described (19).

Denitrification under various NO_3^- concentrations. A randomized complete block experimental design was used, with four treatments and three replicates, and with repeated sampling over time. Treatments were four NO_3^- concentrations (0, 10, 100, and 1,000 mg of potassium nitrate [KNO_3]/liter), and sampling was done at six time points: 0, 2, 4, 6, 8, and 24 h. The experiment was conducted twice. The two experiments were treated as blocks for the analysis of variance.

P. mandelii PD30 was precultured aerobically in TSB medium and shaken in phosphate-buffered saline for 1 h, and the cell biomass was used to inoculate liquid cultures containing 0, 10, 100, or 1,000 mg of KNO_3 /liter. Denitrification conditions were established as previously described by flushing the flask and replacing the headspace atmosphere with helium (19). At each time point, 3-ml samples were collected and assayed for the optical density (OD), RNA isolation, and NO_3^- and NO_2^- analyses. In addition, 12-ml headspace gas samples were obtained to allow for N_2O analysis.

Denitrification in the presence or absence of acetylene. A randomized complete block design was used. The experiment had two treatments and three replicates, and used repeated sampling over time. Treatments consisted of two levels (acetylene+, where 10% of the headspace was composed of acetylene, and acetylene-, where 100% of the headspace was composed of helium), and sampling was done at 2 and 6 h to ensure that the NO_3^- concentration was not limiting. The experiment was conducted once. *P. mandelii* PD30 cultures were established as described above in TSB medium supplemented with 0.1% KNO_3 . Upon induction of denitrification conditions, the flasks were made anaerobic, by making the atmosphere either 100% helium (acetylene-) or 10% (wt/vol) acetylene-90% helium (acetylene+). At each time point, 3-ml samples were collected and then assayed for total RNA isolation and subsequent denitrification gene expression. Headspace gas samples were also collected and analyzed for N_2O .

Design of *P. mandelii nosZ* quantitative PCR primers. *P. mandelii nosZ* sequences (DQ377802, DQ377780, and DQ377774) were aligned to find unique, conserved regions using the MegAlign (Lasergene 7) software (Applied Biosystems, Foster City, CA). Primers were selected based on standard conditions for real-time quantitative PCR using PrimerSelect (Lasergene 7). The specificity of the primers was tested and verified with *P. mandelii* genomic DNA in a PCR. The product was sequenced and submitted to BLAST (NCBI) to ensure specificity (S. Henderson, unpublished data).

Gene expression quantification. The RNeasy minikit (Qiagen, Inc., Mississauga, Ontario, Canada) was used for total RNA isolation and modified as described by Saleh-Lakha et al. (19). RNA was quantified by using Ribogreen

RNA quantitation reagent (Molecular Probes, Eugene, OR). Gene expression quantification was performed on the Bio-Rad iCycler iQ detection system (Bio-Rad Laboratories, Mississauga, Ontario, Canada). Previously designed primers from our research group targeting the *cnorB* region and *nirS* region were used (8). The conditions for one-step quantitative PCR targeting of *P. mandelii cnorB* gene and *nirS* gene were the same as previously described (8).

P. mandelii nirS gene expression was quantified by using one-step quantitative PCR with 50 ng of total RNA template, 12.5 μ l of 2 \times master mix from the Qiagen QuantiTect SYBR green reverse transcription-PCR kit, 300 nM forward primer (5'-ACCGCGCAACAACCTCCAACA-3'), 500 nM reverse primer (5'-CCGCCCTGGCCCTTGAGC-3'), and 0.25 μ l of reverse transcriptase in a final volume of 25 μ l. The thermal cycling conditions were as follows: 30 min at 50°C, 15 min at 95°C, followed by 40 cycles of 15 s at 95°C, 30 s at 68.4°C, 30 s at 72°C, and 15 s at 83°C. The data collection was performed during the last step of each cycle at a temperature of 83°C. In a 40-cycle PCR, a no-template control was undetected.

P. mandelii nosZ gene expression was quantified by using one-step quantitative PCR with 50 ng of total RNA template, 12.5 μ l of 2 \times Qiagen QuantiTect SYBR green reverse transcription-PCR kit, 300 nM forward primer (5'-GGACTAAA AAGATCTGGGAC-3'), 300 nM reverse primer (5'-GTGTCACGTCTCCAC CTTATC-3'), and 0.25 μ l of reverse transcriptase in a final volume of 25 μ l. The thermal cycling conditions were as follows: 30 min at 50°C, 15 min at 95°C, followed by 40 cycles of 15 s at 95°C, 30 s at 56.4°C, 30 s at 72°C, and 15 s at 80°C. The data collection was performed during the last step of each cycle at a temperature of 80°C. In a 40-cycle PCR, a no-template control was undetected.

During each quantitative reverse transcription-PCR run, standard dilutions of digested plasmid carrying a cloned copy of *nosZ* gene were included to allow for gene quantification. *P. mandelii nosZ* primers were used to produce a *nosZ* PCR product that was then cloned in pGEM-TEasy vector according to the manufacturer's instructions (Promega). Plasmid DNA was extracted by using a plasmid minikit (Qiagen, Inc., Mississauga, Ontario, Canada). The plasmid was linearized by digesting with PstI (Roche, Laval, Quebec, Canada) and heat shocking to deactivate the enzymes. The linearized plasmid was quantified by using PicoGreen (Molecular Probes), and the size of the *nosZ* insert was used to calculate the copy number. The curve was linear over a dilution range of 10^{-3} to 10^{-9} and sensitive to at least 10 copies of *nosZ* per reaction. Positive control consisting of *P. mandelii* genomic DNA isolated as outlined in the DNeasy tissue kit (Qiagen) and quantified by Quant-iT PicoGreen dsDNA assay kit (Molecular Probes) was used in each quantitative reverse transcription-PCR run.

NO_3^- , NO_2^- , and N_2O analyses. Frozen culture supernatant samples were analyzed for NO_3^- -N and NO_2^- -N concentrations at the Soil and Nutrient Laboratory (Laboratory Services Division, University of Guelph, Guelph, Ontario, Canada). N_2O analysis of headspace gas was performed by gas chromatography as previously described (8). Calculated N_2O values were corrected for dissolved N_2O in the flask by using the Bunsen absorption coefficient (16) and for changes in pressure and gas concentration attributed to sampling.

Statistical methods. All parameters were tested for normality using the UNIVARIATE function in the SAS System for Windows (version 8; SAS Institute, Inc., Cary, NC), and a log transformation was performed when required. Analysis of variance was performed by using the MIXED procedure of SAS. For the first experiment, the statistical model treated duplicate experiments as blocks in order to pool data from the two experiments, and the REPEATED function was used to account for repeated sampling of flasks over time. Where there was a significant nitrate treatment by time interaction, treatment means were compared by LSMEANS with a Tukey's adjustment. Statistical significance was accepted at $P < 0.05$. Treatment means and standard errors presented in the figures are calculated from nontransformed data.

RESULTS

***P. mandelii* growth and denitrification gene expression at 0, 10, 100, and 1,000 mg of KNO_3 /liter.** Cell density increased over time in all NO_3^- treatments (Fig. 1). At 8 h, the OD at 600 nm for the 1,000-mg N/liter treatment of 0.36 was higher than for the other three NO_3^- treatments (average OD of 0.23). Similarly, at 24 h, *P. mandelii* cell cultures grown at 1,000 mg of N/liter had a higher OD at 600 nm, 1.01, than with all other NO_3^- treatments (average OD of 0.45).

NO_3^- treatment had a significant effect on *nirS* gene expression in *P. mandelii* cultures, and the response differed over




FIG. 1. *P. mandelii* growth in TSB medium supplemented with 0 mg (◆), 10 mg (■), 100 mg (◇), or 1,000 mg (□) of KNO_3 /liter was measured by determining the OD measurements at 600 nm. Error bars indicate ± 1 standard error of the mean (SEM) ($n = 6$).

time. For the 0-mg/liter KNO_3 treatment, *nirS* gene expression increased 12-fold from 0 to 2 h, with 2.7×10^8 transcripts/ μg RNA, followed by a significant decline in *nirS* gene expression at 4 h, and then remained constant thereafter at an average of 3.6×10^7 transcripts/ μg RNA between 4 and 24 h (Fig. 2A). In contrast, *nirS* gene expression increased 143-fold in *P. mandelii* cultures grown at 10 and 100 mg of KNO_3 /liter (average of 6.4×10^8 transcripts/ μg RNA) and 225-fold in *P. mandelii* cultures grown at 1,000 mg of KNO_3 /liter (2.2×10^9 transcripts/ μg RNA) to reach a maximum point at 4 and 6 h, respectively, compared to the start of the incubation. Subsequently, a significant decline of in *nirS* gene expression was measured at 8 and 24 h in *P. mandelii* cultures grown at 100 and 1,000 mg KNO_3 /liter, respectively. There was no significant difference in *nirS* gene expression between 4 and 24 h in *P. mandelii* cultures grown at 10 mg of KNO_3 /liter (average of 1.3×10^8 transcripts/ μg RNA). *nirS* gene expression was not significantly different among treatments at 24 h, with an average of 1.3×10^7 transcripts/ μg RNA.

NO_3^- treatment had a significant effect on *cnorB* gene expression in *P. mandelii* cultures and the response differed over time. Cultures grown at 0 mg of KNO_3 /liter, demonstrated an 86-fold induction in *cnorB* gene expression, where *cnorB* gene transcripts increased from an average of 1.1×10^5 transcripts/ μg RNA at 0 h to 9.2×10^6 transcripts/ μg RNA at 2 h (Fig. 2B). The number of *cnorB* transcripts subsequently declined to 1.6×10^5 transcripts/ μg RNA at 6 h and then did not change until the end of the incubation at 24 h. Maximum *cnorB* gene expression in *P. mandelii* cultures grown at 10, 100, and 1,000 mg of KNO_3 /liter was measured at 4 h, where *P. mandelii* cultures exhibited a 949-fold increase for the 10-mg/liter KNO_3 treatment (1.4×10^8 transcripts/ μg RNA at 4 h) and a 4,550-fold increase for the 100- and 1,000-mg/liter KNO_3 treatments (average of 4.8×10^8 transcripts/ μg RNA at 4 h), compared to the start of the incubation. For *P. mandelii* cultures grown at 10 and 100 mg of KNO_3 /liter, *cnorB* gene transcripts started to decrease after 4 h and reached time zero transcript levels by




FIG. 2. *nirS* (A), *cnorB* (B), and *nosZ* (C) gene expression in *P. mandelii* grown in TSB medium supplemented with 0 mg (◆), 10 mg (■), 100 mg (◇), or 1,000 mg (□) of KNO_3 /liter. Error bars are ± 1 SEM ($n = 6$ for *cnorB* and *nirS* and $n = 3$ for *nosZ*). Values were calculated from the line of best fit described by the linear equation for *cnorB*: $y = -4.112x + 46.661$ ($r^2 = 0.99$) for the first experiment and $y = -4.800x + 53.412$ ($r^2 = 0.99$) for the second experiment. For *nirS*, $y = -2.883x + 45.326$ ($r^2 = 0.99$) for the first experiment and $y = -2.377x + 42.393$ ($r^2 = 0.86$) for the second experiment. For *nosZ*, $y = -4.414x + 46.328$ ($r^2 = 0.99$).

6 h with the 10-mg/liter KNO_3 treatment (5.1×10^5 transcripts/ μg RNA) and by 8 h with the 100-mg/liter KNO_3 treatment (9.2×10^5 transcripts/ μg RNA). Both treatments maintained decreased *cnorB* gene expression at 24 h (average of 4.0×10^5 transcripts/ μg RNA). For *P. mandelii* culture grown at 1,000 mg KNO_3 /liter, *cnorB* gene transcript numbers remained unchanged between 4 and 8 h (average of 2.6×10^8 transcripts/ μg RNA) subsequently decreased to 3.8×10^6 transcripts/ μg RNA at 24 h.

There was no significant effect of treatment or time and no significant treatment-time interaction on *nosZ* gene expression in *P. mandelii* cultures (Fig. 2C). *nosZ* gene expression averaged 3.8×10^4 transcripts/ μg RNA for all treatments over 24 h.

NO_3^- and NO_2^- concentration and denitrification in *P. mandelii* culture grown at 0, 10, 100, and 1,000 mg of KNO_3 /liter. NO_3^- treatment had a significant effect on NO_3^- and NO_2^- concentrations in *P. mandelii* cultures, and the response differed over time. NO_3^- and NO_2^- concentrations were not significantly different among *P. mandelii* cultures grown at 0, 10, and 100 mg KNO_3 /liter (averages of 1.5 mg of NO_3^- -N/liter and 1.3 mg of NO_2^- -N/liter, respectively, from 2 to 8 h (Fig. 3A and B). However, in *P. mandelii* cultures grown at 1,000 mg of KNO_3 /liter, NO_3^- concentrations remained high for 4 h (average of 138 NO_3^- -N/liter), subsequently declined, and were undetectable by 24 h. NO_2^- concentrations, in *P. mandelii* cultures grown at 1,000 mg/liter KNO_3 reached a maximum of 60 mg of NO_2^- -N/liter at 6 h, subsequently declined, and were undetectable by 24 h. The magnitude of NO_3^- loss (77 ppm) was proportional to the increase in NO_2^- (54 ppm) at 6 h, which was the time at which significant differences in both NO_3^- and NO_2^- concentrations in *P. mandelii* cultures grown at 1,000 mg of KNO_3 /liter was observed.

NO_3^- treatment had a significant effect on the cumulative denitrification in *P. mandelii* cultures, and the response differed over time. Denitrification was undetectable in *P. mandelii* cultures grown at 0 mg of KNO_3 /liter and, although detectable with the 10- and 100-mg/liter KNO_3 treatments, was low, with averages of 1.0 and 7.2 μmol at 24 h, respectively (Fig. 3C). However, in *P. mandelii* cultures grown at 1,000 mg of KNO_3 /liter, denitrification was first detected at 4 h and rapidly increased to 98 μmol by 24 h.

Effect of acetylene on *P. mandelii* denitrification gene expression and denitrification activity. There was no significant effect of acetylene treatment or time, or acetylene treatment by time interaction, on *nirS* and *cnorB* gene expression in *P. mandelii* cultures (data not shown). In contrast, acetylene treatment had a significant effect on *nosZ* gene expression in *P. mandelii* cultures, and the response differed over time. The presence of acetylene in the headspace of denitrifying *P. mandelii* cultures inhibited *nosZ* gene expression, with an average of 6.4×10^5 transcripts/ μg RNA in the presence of acetylene and 4.2×10^6 transcripts/ μg RNA in the absence of acetylene between 2 h and 6 h (Fig. 4).

DISCUSSION

This study demonstrated that the higher the concentration of NO_3^- present, the greater the induction and the longer the denitrification genes remained expressed. Maximum *nirS* gene




FIG. 3. NO_3^- (A) and NO_2^- (B) concentrations and denitrification (C) in TSB medium supplemented with 0 mg (◆), 10 mg (■), 100 mg (◇), or 1,000 mg (□) of KNO_3 /liter. Error bars are ± 1 SEM ($n = 6$).

expression was 12-fold at 2 h for cultures grown at 0 mg of KNO_3 /liter, 43-fold at 4 h in cultures grown at 10 and 100 mg of KNO_3 /liter and 225-fold at 6 h in cultures grown at 1,000 mg of KNO_3 /liter. Similarly, *cnorB* gene expression reached a maximum with an 86-fold increase at 2 h in cultures grown at 0 mg of KNO_3 /liter, a 949-fold increase at 4 h for the 10-mg/liter KNO_3 treatment, and a 4,550-fold increase at 4 h for the 100




FIG. 4. *nosZ* gene expression in *P. mandelii* grown in TSB medium supplemented with 1,000 mg of KNO_3 /liter, grown in the presence (■) or absence (□) of acetylene. Error bars indicate ± 1 SEM ($n = 6$ for *cnorB* and *nirS* and $n = 3$ for *nosZ*). Values were calculated from the line of best fit described by the linear equation for *cnorB*: $y = -3.275x + 35.640$ ($r^2 = 0.98$) for the first experiment and $y = -2.772x + 34.347$ ($r^2 = 0.99$) for the second experiment. For *nirS*, $y = -2.648x + 46.202$ ($r^2 = 0.99$) for the first experiment and $y = -3.014x + 47.619$ ($r^2 = 0.99$) for the second experiment. For *nosZ*, $y = -3.799x + 44.825$ ($r^2 = 0.99$).

and 1,000 mg/liter KNO_3 treatments. Increased *cnorB* gene expression was also measured between 1 and 6 h in a study by Saleh-Lakha et al. (19) in *P. mandelii* cultures grown in the presence of 1,000 mg of KNO_3 /liter under anaerobic conditions. Hartig and Zumft (10) also measured *nirS* and *norCB* gene expression via Northern blot analysis and reported constant gene expression for a 3-h observation period in denitrifying cultures of *Pseudomonas stutzeri* grown in the presence of 1,000 mg of KNO_3 /liter. The present study demonstrated that continued transcription of *nirS* and *cnorB* genes in *P. mandelii* required the presence of NO_3^- .

In our study, *nosZ* gene expression was not affected by various NO_3^- concentrations, nor was the typical pattern of gene induction and gene expression observed, as was the case for *nirS* and *cnorB* genes. There was no *nosZ* gene induction, and the number of *nosZ* transcripts measured was about 10^4 - and 10^3 -fold lower than both *nirS* and *cnorB* gene expression, respectively. This observation is contrary to previous studies where similar induction patterns and coordinated expression of *nirS*, *cnorB*, and *nosZ* genes was observed in *P. stutzeri* (10, 13). The difference in *nosZ* gene expression between the present study and previous studies could be attributed to the use of acetylene. Acetylene blockage of the nitrous oxide reductase results in N_2O accumulation, which in turn, could affect *nosZ* regulation. It is also possible that acetylene affects other metabolic processes, which indirectly affects *nosZ* gene regulation or expression. Alternatively, *nosZ* gene expression could also have been affected by differences in the experimental conditions used to establish precultures and denitrification conditions, or alternatively, a different mode of *nosZ* gene regulation in *P. mandelii*.

We also investigated the effect of variable NO_3^- concentrations on denitrification intermediates including, specifically,

NO_2^- . Although insignificant NO_2^- concentrations were observed under limiting (0-, 10-, and 100-mg/liter KNO_3) NO_3^- treatments, 60 mg of NO_2^- -N/liter accumulated at 6 h in *P. mandelii* cultures grown with 1,000 mg of KNO_3 /liter. There are very few studies that have measured NO_2^- accumulation in the medium of denitrifying cultures. Carlson and Ingraham (5) compared the accumulation of denitrification intermediates in *P. stutzeri*, *P. aeruginosa*, and *P. denitrificans* and concluded that all three species accumulated NO_2^- at different rates, with *P. stutzeri* having the highest NO_2^- accumulation, with an average of 220 μmol , in the presence of 23 mM sodium nitrate. Our previous study also measured insignificant NO_2^- in *P. mandelii* cultures grown at 0 mg of KNO_3 /liter; however, only 8.5 mg of NO_2^- -N/liter accumulated at 6 h in *P. mandelii* culture grown with 1,000 mg of KNO_3 /liter (19). The difference in NO_2^- accumulation in our previous study and the current one can be explained by slightly different growth conditions (larger flask size and culture volume), suggesting that culture conditions can alter metabolism in *P. mandelii* cells and thus affect NO_2^- accumulation.

The effect of NO_3^- concentrations on denitrification activity, as measured by N_2O emissions in the presence of C_2H_2 , demonstrated significant denitrification activity (98 μmol at 24 h) in *P. mandelii* culture grown with 1,000 mg of KNO_3 /liter. A recent study also measured 101 μmol of N_2O at 24 h in a *P. mandelii* culture grown with 1,000 mg of KNO_3 /liter; however, the effects of lower NO_3^- concentrations on denitrification activity were not investigated (19). Carlson and Ingraham (5) measured different accumulation rates of denitrification intermediates (NO_2^-) and products (N_2O and N_2) under unlimited nitrate conditions in pure cultures of *P. stutzeri*, *P. aeruginosa*, and *P. denitrificans* and also concluded that denitrification activity was high under conditions where nitrate was in excess. Our results from the present study demonstrated insignificant denitrification activity under limiting (0-, 10-, and 100-mg/liter) KNO_3^- treatments, suggesting that KNO_3^- concentrations of up to 100 mg/liter were not sufficient to allow for significant denitrification activity. There are no other pure culture studies, to our knowledge, that investigated the effect of limited NO_3^- on denitrification activity. The results presented here demonstrate that *nirS* and *cnorB* responded to nitrate concentrations; however, significant denitrification activity was only observed in culture with 1,000 mg of KNO_3 /liter, indicating that there was no relationship between gene expression and denitrification activity under the conditions tested.

Numerous studies have used acetylene, both in pure culture (3, 19, 25) and soil microcosms (6, 15, 17, 18, 23), as a measure of denitrification. In the present study, denitrification gene expression (*nirS*, *cnorB*, and *nosZ*) was measured in the presence or absence of acetylene in order to determine whether acetylene influences gene expression. Acetylene did not affect *nirS* or *cnorB* gene expression in *P. mandelii*; however, it significantly reduced *nosZ* gene expression. It has been well established that acetylene inhibits the nitrous oxide reductase enzyme resulting in N_2O accumulation (3, 25); however, the effect of acetylene on *nosZ* gene expression had not yet been determined. The present study provides direct evidence that acetylene decreases *nosZ* gene expression. This could be due to an effect of acetylene on general metabolism, which in turn, indirectly affects *nosZ* gene expression. It is also possible that

the N₂O atmosphere in the flask might interfere with *nosZ* gene regulation in *P. mandelii*, although a recent study investigating transcriptional regulation of the *nos* genes for nitrous oxide reductase in *P. aeruginosa* concluded that N₂O did not affect induction of the promoter for *nosZ* (2).

In conclusion, we investigated the effect of various NO₃⁻ concentrations on growth, denitrification gene expression, and cumulative denitrification. The higher the amount of NO₃⁻ present, the greater the induction and the longer the denitrification genes remained expressed. *nosZ* gene expression did not respond to NO₃⁻ treatment under the conditions used. Acetylene negatively influenced *nosZ* gene expression but did not affect *nirS* or *cnorB* gene expression. This observation has important implications in experimental design, especially when investigating *nosZ* gene expression. This research provided evidence for the effect of varying the NO₃⁻ concentration on *nirS* and *cnorB* gene expression; however, significant denitrification activity was only observed in culture with 1,000 mg of KNO₃/liter, indicating that there was no relationship between gene expression and denitrification activity under the conditions tested.

ACKNOWLEDGMENTS

We are grateful to Drucie Janes and Jan Zeng for providing technical support. We are also grateful to Stephen Bowley for providing guidance in the statistical analysis of data.

Funding for this project was supplied by the GAPS program of Agriculture and Agri-Food Canada and an NSERC Strategic Team Grant. S.S.-L. was the recipient of an NSERC doctoral scholarship. S.S.-L. and K.E.S. were also recipients of Ontario Graduate Scholarships. Infrastructure and equipment grants from the Canadian Foundation Innovation and the Ontario Innovation Trust are sincerely acknowledged by J.T.T.

REFERENCES

1. Arai, H., T. Kodama, and Y. Igarashi. 1999. Effect of nitrogen oxides on expression of the *nir* and *nor* genes for denitrification in *Pseudomonas aeruginosa*. *FEMS Microbiol. Lett.* **170**:19–24.
2. Arai, H., M. Masayuki, and Y. Igarashi. 2003. Transcriptional regulation of the *nos* genes or nitrous oxide reductase in *Pseudomonas aeruginosa*. *Microbiology* **149**:29–36.
3. Balderstone, W. L., B. Sherr, and W. J. Payne. 1976. Blockage by acetylene of nitrous oxide reduction in *Pseudomonas perfectomarinus*. *Appl. Environ. Microbiol.* **31**:504–508.
4. Bergaust, L., J. Shapleigh, Å. Frostegård, and L. Bakken. 2008. Transcription and activities of NO_x reductases in *Agrobacterium tumefaciens*: the influence of nitrate, nitrite, and oxygen availability. *Environ. Microbiol.* **10**:3070–3081.
5. Carlson, C. A., and J. L. Ingraham. 1983. Comparison of denitrification by *Pseudomonas stutzeri*, *Pseudomonas aeruginosa*, and *Paracoccus denitrificans*. *Appl. Environ. Microbiol.* **45**:1247–1253.
6. Dandie, C. E., D. L. Burton, B. J. Zebarth, S. L. Henderson, J. T. Trevors, and C. Goyer. 2008. Changes in bacterial denitrifier community abundance over time in an agricultural field and their relationship with denitrification activity. *Appl. Environ. Microbiol.* **74**:5997–6005.
7. Dandie, C. E., D. L. Burton, B. J. Zebarth, J. T. Trevors, and C. Goyer. 2007. Analysis of denitrification genes and comparison of *nosZ*, *cnorB*, and 16S rDNA from culturable denitrifying bacteria in potato cropping systems. *Syst. Appl. Microbiol.* **30**:128–138.
8. Dandie, C. E., M. N. Miller, D. L. Burton, B. J. Zebarth, J. T. Trevors, and C. Goyer. 2007. Nitric oxide reductase-targeted real-time PCR quantification of denitrifier populations in soil. *Appl. Environ. Microbiol.* **73**:4250–4258.
9. Hartig, E., U. Schiek, K. U. Vollack, and W. G. Zumft. 1999. Nitrate and nitrite control of respiratory nitrate reduction in denitrifying *Pseudomonas stutzeri* by a two-component regulatory system homologous to NarXL of *Escherichia coli*. *J. Bacteriol.* **181**:3658–3665.
10. Hartig, E., and W. G. Zumft. 1999. Kinetics of *nirS* expression (cytochrome cd1 nitrite reductase) in *Pseudomonas stutzeri* during the transition from aerobic respiration to denitrification: evidence for a denitrification-specific nitrate- and nitrite-responsive regulatory system. *J. Bacteriol.* **181**:161–166.
11. Jarvis, S. C., D. J. Hatch, and R. D. Lovell. 2001. An improved soil core incubation methods for the field measurement of denitrification and net mineralization using acetylene inhibition. *Nutr. Cycl. Agroecosys.* **59**:219–225.
12. Knowles, R. 1982. Denitrification. *Microbiol. Rev.* **46**:43–70.
13. Körner, H. 1993. Anaerobic expression of nitric oxide reductase from denitrifier *Pseudomonas stutzeri*. *Arch. Microbiol.* **159**:410–416.
14. Körner, H., and W. G. Zumft. 1989. Expression of denitrification enzymes in response to the dissolved oxygen level and respiratory substrate in continuous culture of *Pseudomonas stutzeri*. *Appl. Environ. Microbiol.* **55**:1670–1676.
15. Miller, M. N., B. J. Zebarth, C. E. Dandie, D. L. Burton, C. Goyer, and J. T. Trevors. 2008. Crop residue influence on denitrification, N₂O emissions and denitrifier community abundance in soil. *Soil Biol. Biochem.* **40**:2553–2562.
16. Moraghan, J. T., and R. Buresh. 1977. Correction for dissolved nitrous oxide in nitrogen studies. *Soil Sci. Soc. Am. J.* **41**:1201–1202.
17. Müller, M. M., V. Sundman, and J. Skujins. 1980. Denitrification in low pH spodosols and peats determined with the acetylene inhibition method. *Appl. Environ. Microbiol.* **40**:235–239.
18. Parkin, T. B., A. J. Sextone, and J. M. Tiedje. 1985. Adaptation of denitrifying population to low soil pH. *Appl. Environ. Microbiol.* **49**:1053–1056.
19. Saleh-Lakha, S., K. E. Shannon, C. Goyer, J. T. Trevors, B. J. Zebarth, and D. L. Burton. 2008. Nitric oxide reductase gene expression and nitrous oxide production in nitrate-grown *Pseudomonas mandelii*. *Appl. Environ. Microbiol.* **74**:6876–6879.
20. Smith, M. S., M. K. Firestone, and J. M. Tiedje. 1978. The acetylene inhibition method for short-term measurement of soil denitrification and its evaluation using nitrogen-13. *Soil Sci. Soc. Am. J.* **42**:611–615.
21. Tate, R. L., III. 2000. Denitrification, p. 404–432. *In* Soil microbiology, 2nd ed. John Wiley, Inc., New York, NY.
22. Thomas, K. L., D. Lloyd, and L. Boddy. 1994. Effects of oxygen, pH and nitrate concentration on denitrification by *Pseudomonas* species. *FEMS Microbiol. Lett.* **118**:181–186.
23. Van Raalte, C. D., and D. G. Patriquin. 1979. Use of the acetylene blockage technique for assaying denitrification in a salt marsh. *Mar. Biol.* **52**:315–320.
24. Vollack, K. U., and W. G. Zumft. 2001. Nitric oxide signaling and transcriptional control of denitrification genes in *Pseudomonas stutzeri*. *J. Bacteriol.* **183**:2516–2526.
25. Yoshinari, T., and R. Knowles. 1976. Acetylene inhibition of nitrous oxide reduction by denitrifying bacteria. *Biochem. Biophys. Res. Commun.* **69**:705–710.
26. Zumft, W. G. 1997. Cell biology and molecular basis of denitrification. *Microbiol. Mol. Biol. Rev.* **61**:533–616.