

Field Trip 401, Halifax '03

FIELD TRIP GUIDEBOOK


Tidal Sedimentation in the Cornwallis River - Avon River Estuaries, Bay of Fundy, Nova Scotia: Sedimentology, Morphology, and Natural History

(Text modified from Dalrymple, Amos, and Yeo, 1992)

Ian S. Spooner¹ and G. Yeo²

¹Department of Geology, Acadia University
Wolfville, Nova Scotia, Canada, B4P 2R6

²Saskatchewan Industry and Resources, 2101 Scarth St.
Regina, Saskatchewan, Canada, S4P 3V7



**Northeastern Section, The Geological Society of America
Atlantic Geoscience Society
Joint Meeting
Halifax, 2003**

with

**Eastern Section American Association of Petroleum Geologists (AAPG)
Council on Undergraduate Research (CUR), Geoscience Division
Society of Economic Geologists (SEG)**

© Atlantic Geoscience Society

**Department of Earth Sciences
Dalhousie University
Halifax, Nova Scotia, Canada
B3H 3J5**

**ISBN 0-9683988-5-5
AGS Special Publication Number 17**

**Tidal Sedimentation in the Cornwallis River – Avon River Estuaries,
Bay of Fundy, Nova Scotia:
Sedimentology, Morphology, and Natural History**

Ian S. Spooner¹ and G. Yeo²

¹*Department of Geology, Acadia University, Wolfville, N.S. B4P 2R6*

²*Saskatchewan Industry and Resources 2101 Scarth St., Regina, Saskatchewan, S4P 3V7, Canada
(Text modified from Dalrymple, Amos, and Yeo, 1992*)*

Summary: The Bay of Fundy is an elongate, structurally-controlled basin situated on the east coast of Canada between the provinces of New Brunswick and Nova Scotia. It is widely known and frequently cited for its very large tidal range, which averages 12.0 m. Most of the floor of the outer Bay of Fundy is presently non-depositional or erosional as a result of the strong tidal currents and the minimal input of sediment from terrestrial sources. By contrast, the two major estuaries at the head of the Bay (the northern Chignecto Bay arm and the eastern Minas Basin-Cobequid Bay arm) are experiencing rapid accumulation of sediment. In these two arms Holocene tidal deposits are in places more than 10m thick.

This field trip will focus on the sedimentology, morphology and natural history of the Windsor Bay area with emphasis on the Cornwallis and Avon River Estuaries. This region is one of the more thoroughly studied macrotidal areas in the world sedimentologically. This work has covered virtually all aspects of the sedimentation processes and deposit characteristics. As a result, this field trip will focus on an overview of tidal deposits and the response of the system to changing sea levels and tidal ranges since deglaciation. We will also visit some historic sites in the region and explore how the natural environment has influenced human activity in the region.

**Disclaimer: This guide book contains, in part, an abbreviation of material originally presented in the Geological Association of Canada-Mineralogical Association of Canada (GAC-MAC) Field Excursion A-8: Guidebook (1992) which is available through many library and private collections. It can also be obtained by contacting:*

*Atlantic Geoscience Society
c/o The Department of Earth Sciences,
Dalhousie University, Halifax Nova Scotia, Canada B3H 3J5*

Please refer to the above volume for a much more thorough treatment of many of the topics discussed herein. Access information to many of the sites can also be found in the 1992 publication.

WARNING TO GUIDEBOOK USERS- SAFETY PRECAUTIONS

THE EXTREMELY RAPID RISE OF WATER LEVELS ON THE FLOODING TIDE (>2 METRES/HOUR!), COUPLED WITH THE SOFT, STICKY MUD, CREATES POTENTIALLY DANGEROUS SITUATIONS.

The locals also tell stories of people who have become stuck in the mud and drowned. Consequently, CAUTION MUST BE EXERCISED AT ALL TIMES. Please stay with the group and listen to the leader's instructions. Do not wander off on your own!

With respect to the mud, the rule of thumb is that if you sink in deeper than mid-calf, return to solid ground as quickly as possible. DO NOT STOP MOVING as this will allow the mud to "freeze", trapping you more firmly.

THE AUTHORS ACCEPT NO RESPONSIBILITY FOR INJURIES OR DEATH SUSTAINED BY ANY USER OF THE GUIDEBOOK.

INTRODUCTION: *SEDIMENTOLOGY OF THE CORNWALLIS RIVER ESTUARY*

Surficial Geology and Geomorphology

The Cornwallis River estuary lies at the southwestern corner of the Minas Basin, and is the focus of drainage from the eastern part of the Annapolis Valley (Fig. 1). The Annapolis Valley is bounded on the north by the North Mountain cuesta that is capped by Triassic-Jurassic basalt, and on the south by a highland area called the South Mountain that is underlain by Paleozoic metasedimentary and granitic rocks. The Valley floor is underlain by red fluvial sandstones of the Triassic Wolfville Formation, and by the conformably overlying fluvial and lacustrine mudstones of the Blomidon Formation (Fig. 2a). The Wolfville sandstone is exposed in low cliffs along the shore of the Cornwallis River estuary. Unconsolidated, surficial sediments include red silty till, outwash sands and muds, and the recent tidal deposits.

The Cornwallis River estuary is a convergent (trumpet-shaped) estuary (Fig. 3). Its mouth lies between Kingsport and Long Island-Evangeline Beach, while the head of tide is about half a kilometre upriver from Kentville. It is strongly macrotidal, having a spring tide range in the outer estuary of about 15 m, and a neap tide range of approximately 9 m (Fig. 2b). Average salinity decreases headward, from about 29‰ at the mouth to less than 0.1 ‰ at the Kentville Bridge. In the outer estuary, higher salinities are observed along the south shore, where the flood tide is strongest (Daborn and Pennachetti, 1979). The total volume of water in the estuary, as estimated from bathymetry (Yeo; in Daborn, 1991), is about $138 \times 10^6 \text{ m}^3$. The mixing index (the volumetric ratio of fresh water input during a half-tidal period to marine water entering during a flood tide; Schubel, 1971) is estimated to be 0.004 (Dickinson, 1991). This is well within the range for well-mixed, Type C (Pritchard in Schubel, 1971) estuaries.

The Cornwallis estuary is dominated by muddy sediments (Fig. 4). The outer estuary (i.e., seaward of Wolfville; Figs. 3, 4), in particular, is characterized by broad mudflats instead of the erosional foreshores and elongate sand bars which are so abundant in Cobequid Bay (Fig. 1). The chief reason for the difference is the nature of the sediment source; because of the extensive outcrops of muddy Blomidon Formation at Cape Blomidon to the north, the Cornwallis estuary receives little sand. The abundance of mud in turn causes this area to be more productive biologically than Cobequid Bay, and organisms have a much greater influence on sedimentation and erosion, especially in the outer estuary. Partly because of the abundant mud, the sedimentology of the Cornwallis estuary has been little studied until recently.

The Inner Cornwallis Estuary

The Cornwallis River is tidal to a point about 0.5 kilometre upriver from Kentville. The channel from Wolfville to the head of tide is relatively narrow, has steep banks, and lacks the extensive tidal flats characteristic of the outer estuary. The width to depth ratio is only on the order of 10-20 through the first 10-15 km below the tidal limit, and then increases dramatically beyond this. The bankfull capacity of the inner estuary above the Port Williams bridge (Stop 4) is about $8 \times 10^6 \text{ m}^3$ of which 97.5% is tidal. Hence, average tidal discharge is about 360 cubic metres per second.

Suspended sediment concentration in the inner estuary ranges up to 5 g/l (Daborn and Pennachetti, 1979) with a turbidity maximum located in the vicinity of the town of New Minas. In contrast suspended sediment concentration at mouth of the outer estuary is about 10 mg/l or less. Salinity ranges from 0 ‰ at Kentville to 30 ‰ at Wolfville, while water temperature is variable. Due to the strong currents, high turbidity, low average salinity, and variable temperature within the inner estuary, very few organisms live here except in the salt marshes. This is in marked contrast




Figure 1. Map of the Minas Basin-Cobequid Bay-Salmon River area showing the location of the stops to be visited on this trip




Figure 2(a). A: Location map of eastern Canada B: Geological and location map of the Bay of Fundy region. M.B.- Minas Basin; C.B. Chignecto Bay.





Figure 2(b). History of sea level and tidal range in the inner Bay of Fundy, based on data from Chignecto Bay (after Amos and Zaitlin, 1985). A similar pattern of sea-level and tidal range oscillations have occurred in the Minas Basin but with smaller amplitudes.

to the high biological productivity of the outer estuary. Benthic diatoms and colloform bacteria are abundant however (Dickinson, 1991); faecal colloform numbers, for example, are typically above 200 per ml (except in the winter!).

The Land of Evangeline

Throughout the field trip we will be talking about the natural history of the Minas Basin Region. In particular we will be visiting Grand Pré National Park and the Grand Pré dykelands which consist of over 8000 ha of farmland below sea level. The story of the Grand Pré Acadian's (greatly condensed!) is as follows. In 1680 Acadians first moved to the region from Port Royal and began reclaiming saltmarsh from the tides (see **Farming** below). As the French population of Grand Pré increased and spread over a wide district, the Acadian population exceeded that of the English at Annapolis. Gradually they were drawn into the conflict waged between the French (who controlled Louisbourg in Cape Breton Island and Quebec, and the English who controlled much of the Bay of Fundy region (including Grand Pré) and New England.



Evangeline Beach - Grand Pré area with Minas Basin and Blomidon in the Background

The colonial power struggle between France and England was coming to a climax. Between 1755 and 1763, British authorities enforced a deportation order. This traumatic event became known as the Expulsion. This event has been immortalized by the American poet Henry Wadsworth Longfellow, when he wrote his epic poem "Evangeline".

Acadian Farming

There was little that influenced the development of the Acadian way of life quite as much as their method of farming. Fortunately, among the people who settled in the Annapolis Basin area, there were some who were already familiar with methods of dyking practiced in France, and they recognized the agricultural potential of the tidal salt marshes. The new settlers moved quickly to build dykes along the outer marsh areas. Sometimes these dykes were built by driving five or six rows of logs into the ground, laying other logs one on top of the other between these rows, filling all the spaces between the logs with well packed clay and then covering everything over with sods




Figure 3. Bathymetry of the outer Cornwallis River. Datum is approximately mean low water. Note the development of flood "barbs" along the west side of the Cornwallis channel.




Figure 4. Facies distribution in the outer Cornwallis River estuary. Note the sandy facies are very restricted in extent, while fringing mudflat facies are extensive.

cut from the marsh itself. Sometimes dykes were built by simply laying marsh sods over mounds of earth. The Acadians devised a system of drainage ditches combined with an ingenious one-way water gate called an aboiteau. The aboiteau was a hinged valve in the dyke which allowed fresh water to run off the marshes at low tide but which prevented salt water from flowing onto the dyked farmland as the tide rose. After letting snow and rain wash away the salt from the marshes for between two and four years the Acadians were left with fertile soil which yielded abundant crops.

Field Trip Stops

Please note that some of these stops may be omitted due to road/access conditions!

Stop 1: Windsor Causeway Mudflat (Fig. 1; Stop 1):

Windsor causeway (Figure 1; Stop 1) is a solid-fill barrier across the Avon River estuary which was built in 1970 to support Highway 101 and a rail link between Halifax and the Annapolis Valley. The causeway blocked tidal flow into the inner Avon River estuary, thereby reducing the tidal prism by $4.2 \times 10^6 \text{ m}^3$. This resulted in a significant reduction in tidal currents near the causeway, an effect that decreased exponentially with distance down estuary. Prior to construction, this part of the estuary had sandy mudflats and marshes along its margins. Ships once docked at Windsor against the crib work that now overlooks the permanently inundated river above the causeway. The region upriver of the causeway was converted from an inner estuarine environment where fines accumulated in a strongly oscillating flow, to a fluvial/lacustrine environment with flow in a seaward direction only. Seawards of the causeway, rapid deposition of muddy sediments and a dramatic shallowing resulted from the reduction in tidal current speeds. Welded to the causeway we see a region of lush salt marsh that gives way seawards to a mud flat partially colonized by *Spartina* sp. The mudflat falls away to a silt-dominated reach near the outfall of the St. Croix River and thereafter to the upper- flow-regime and rippled sand flats .

The mudflat in front of the causeway has accumulated 7m over the last 21 years. The majority of this accumulation took place during the first 5 years after construction. The sediments at that time were high in water content, low in bulk density, and of very low bearing capacity. The infauna were restricted to a few species only (mainly a bulbous form of the bivalve *Macoma balthica*). Winter freezing of the mudflat occurred due to the high water content, and large blocks of ice and sediment were commonly littered across the flats. The blocks, which reached 5 m in diameter, were periodically lifted on high spring tides and redistributed throughout the estuary, carrying large volumes of fines with them. In the process, the level of the mudflat was lowered by 2-3 m depending on the severity of the winter. Accumulation recurred during the subsequent spring and summer. The mudflats remained in this dynamic balance until approximately 1985. We presume that throughout this period the mudflat was becoming progressively consolidated by normal geostatic and ice loading. In 1985, the first evidence of salt marsh colonization appeared. Since 1985 the salt marsh has spread dramatically to cover much of the southeastern part of the flat, the region where the highest rates of initial accretion were measured.

Aerial photography between 1973 and 1979 shows that the mudflat has prograded approximately 5 km seawards of the causeway, to partially cover previously-existing sand bars. The remaining sand bars in the inner estuary also show evidence of long-term accretion. A comparison of bathymetric data collected in 1969 and 1976 show that 1-2 m of accretion occurred in that time interval. It is not possible to discriminate natural versus man-made effects in this case. Nevertheless, the estuary appears to be importing sand and fines, and filling throughout its length.

Stop 2 (Optional): Penny Beach: Coastal Erosion – Avon River Estuary (Fig. 1; Stop 2).

At this site (Fig. 1: Stop 2) you will be exposed to excellent examples of coastal erosion and local strategies to mitigate the process. Coastal erosion in the Minas Basin can be as much as 30 cm/year, locally rates of > 1m/year have been noted. Most erosion occurs during the winter months as storms are frequent and often are accompanied by northwesterly winds allowing waves to build up over a maximum fetch of 30 kms. Storms that coincide with high and/or spring tides are especially damaging. As well, both chemical and mechanical (freeze-thaw) weathering can greatly accelerate coastal erosion. Local strategies for dealing with erosion are varied, most property owners use either North Mountain Basalt or South Mountain granite as armour. Some have used (with varying degrees of success) coarse fill and brush. In many cases the placement of armour has exacerbated the problem or, worse yet, transferred the problem down the coastline.

Avon River Estuary: The Southern Bight

The channel of the Avon River Estuary (also called the Southern Bight) consists of a rock-bound, wave-cut shelf no deeper than 10 m below lower low water (LLW). This platform is incised by a paleo-drainage system that is graded to approximately 50 m below LLW. Elongate tidal sand bars have the form of ebb-tidal deltas and are covered by active sand waves and megaripples are found in this channel. By far the greatest amount of sand in the Minas (Windsor) Basin is supplied by erosion of the Triassic sandstone cliffs that border the basin. Most of the sand that is released is either trapped in Cobequid Bay, or moves westwards along the south shore of the Minas Basin and enters the Avon River estuary. These flats at the outer end of the estuary are non-cohesive, silty mudflats that lack large-scale bedforms. They are, however, extremely productive biologically.

Penny Beach:

Penny Beach is in a transgressive phase, migrating landward over saltmarsh peat that is up to 2.5 m thick and is accumulating in the basin landward of the present beach. This basin was formerly more extensive and the peat underlies the present sand barrier., occasionally cropping out on the beach face.

Stop 3: Grand Pré National Park, Evangeline Beach (The Guzzle Section, Fig. 1; Stop 3)

This area, which is known as Grand Pré ("large meadows") is best known as the site of a large Acadian settlement, also known as Les Mines (Fig. 1; Stop 3). This was derived from the name given to this arm of the Bay of Fundy by Champlain on the basis of his copper discovery at Cap D'Or west of Parrsboro in 1607 (promotion of the mineral potential of Nova Scotia is nothing new!) Acadian settlers inhabited the area from the late 1600's to 1755, when they were deported to Louisiana by the British. The ships carrying the Acadians set sail from a tidal creek a short distance to the east. Very little remains from this part of the colonial era, except for the field patterns which follow the early dyke systems. Most of the area between the railway line at the foot of the Wolfville Ridge and Long Island was formerly salt marsh that lay below the highest high tide level. It was originally reclaimed for agricultural use by the Acadians.





Figure 5. Sketch map of The Guzzle area at Evangeline Beach showing the distribution of major facies and features.



A wooden aboiteau (one way drainage valve) in a dyke near Delhaven

Looking northeast across tidal flats one can see Boot Island, a parcel of land separated from the mainland by a tidal gully called The Guzzle. Boot Island was joined to the mainland during the Acadian occupation, and has since become isolated by the headward erosion of a tidal creek used in the deportation of the Acadians. According to a local tradition, the erosion developed from a ditch dug by a farmer. Successive maps show that the channel remained relatively narrow until the early part of this century. Between 1910 and 1940 erosion accelerated greatly but there has been relatively little erosion since then (Bleakney and Davis, 1983).

At the high water line we find a thin, modern salt marsh deposit (Fig. 4) that is up to 2 m thick. Three older rooted horizons between 8 and 75 cm thick may be visible, alternating with grey silty clay layers about 20 cm thick (Ferguson, 1983). These sediments rest on a compact red boulder till, whose matrix is very poorly sorted with a mean size in the coarse silt range. Both units are being eroded by wave activity as the local transgression continues. The product of this erosion is a high tide beach made of sandy gravel and cobbles that grades seawards into a thin mudflat. Cobbles and boulders that appear to be derived from the glacial till litter the intertidal zone.


Approximately 200 m seaward of the salt marsh along the west bank of The Guzzle you will encounter a series of tree stumps in growth position in a forest peat bed (Fig. 5). These mark the former presence of a pine-cedar-hemlock-oak forest (Goldthwaite, 1924) that is dated at 4470 to 3100 years B.P. (Harrison and Lyon, 1963; Bleakney and Davis, 1983). This fossil forest, which is present throughout the southern bight of the Minas Basin, provides graphic evidence of the sea-level lowstand and subsequent rise (Fig. 2b). The trees grew on exposed glacial till in areas near the present shore, and on grey and red laminated mud in more seaward areas. These muds, which also outcrop in other places in The Guzzle, may date from the period of high sea level that accompanied deglaciation 14,000 yr BP (Fig. 2b). A relict salt marsh peat overlies the forest soil layer (Fig. 5).

Rising sea level is indicated by the occurrence of the subtidal oysters (Fig. 5) above salt-marsh deposits and the fossil forest. Following the deposition of the oysters and the overlying mud layer, mean sea level and the high tide level have continued to rise to their present elevations. The high tide level in particular must have risen nearly 10 m as the oysters are situated approximately 13 m below the present-day high water level. By contrast, the low water level has remained

approximately constant, or has fallen slightly, as the oysters are located only slightly above the spring low tide elevation.

A number of cobble-boulder ridges which trend approximately parallel to the modern shoreline are present in the lower half of the intertidal zone. Encrustation of the rocks by large barnacles indicates that these ridges are not active deposits. They are interpreted as either relict low tide, boulder barricades (Dalrymple *et al.*, 1982), or as shoreline ice-push features. Rows of wooden stakes in this area are remnants of old fishing weirs (Fig. 5)

An elongate sand bar is developed in the Guzzle (Figs. 4, 5). This sand bar is asymmetric, rising gently towards its eastern side near the main channel. The sand is well-sorted and medium grained, and small and medium dunes are developed locally. Flood flow dominates the northern part of the bar while ebb flow dominates the southern part (McDow, 1991). The sand bar crest represents an area of zero net bedload transport (i.e., ebb and flood transport are in balance). The overall morphology and distribution of large-scale bedforms suggests that sand transport is counterclockwise around the bar. Possible sources for the sand include outwash deposits such as those on Boot Island and along the lower Gaspereau River, or the sand bars at the mouth of the Avon River. Comparison of 1963 and 1987 airphotos shows that the bar has changed little, hence little or no sand is presently being supplied to it.



Dune with superimposed ripples at The Guzzle

Lunch!!

Stop 4: Port Williams Bridge : Tidal Sand Bars (Fig. 1; Stop 4)

CAUTION: THIS IS A VERY BUSY ROAD. WATCH FOR TRAFFIC!

The Port Williams bridge locality (Fig. 1; Stop 4) lies at a transition from a relatively straight channel with bank-attached bars, and a more sinuous, meandering channel (Fig. 6). The well-




Figure 6. Sketch map of the Cornwallis River at the Port Williams bridge.

developed tidal point bar that is present downstream of the bridge is the focus of this stop. Other bars located further upstream are described in subsequent paragraphs.

Port Williams Bar (Downstream Bar)

The channel here is U-shaped, with steep banks and a relatively flat bottom. At low water, the channel width is about 42 m and depth is 0.5 m, but this varies with fluvial discharge. On a high spring tide, the water depth is 12 m and the width is about 130 m. The difference between the cross-sectional areas of the channel at spring high tide (1169 m²) and low water (20 m²) indicates that the fluvial discharge is less than 2% of total flow here. Suspended sediment concentration, salinity and temperature generally increase on the flood and decrease on the ebb. Except at slack water, the flow is stratified, with the stratification being strongest during the first two hours of flood tide. The average salinity observed at the Port Williams bridge was 21‰, and the maximum suspended sediment concentration was 442 mg/l

Muddy suspension deposits accumulate on the bank at high water, but undercutting of the oversteepened banks results in slumping. Here the banks are stabilized artificially by rip-rap, to prevent erosion of the dykes. As observed elsewhere in the Cornwallis estuary, there is essentially no long-term net accretion or erosion apparent at the Port Williams bar. The lower portion of the point bar (Fig. 6) is composed predominantly of sandy bedload deposits, although organic debris is noticeably more abundant at the upriver end. Ebb-dominant bedforms and structures are preserved on the upriver end of the bar, whereas flood-dominant structures are present on the downriver end because that area is sheltered from the full force of the ebb currents.



The Port Williams Bar downstream of the Port Williams bridge


The largest bedforms occur on the lower part of the bar, near the centre of the channel, and consist of medium dunes of two types. Along the margin of the low-tide channel at the upriver end of the bar, sinuous-crested dunes with avalanche-type foresets and current-lineated topsets may form a point projecting into the channel. These are composed of very coarse sand and granules, and have wavelengths of 2-6 m and heights of about 0.5 m. A box core from the sediments below the dunes in this area revealed preserved, ebb-dominant, planar cross beds in sets only 4-6 cm thick. No muddy drapes were present. Higher on the bar in the same area, dunes with rounded, relatively straight crests may be present in the medium-grained, moderately sorted sand that characterizes this topographically higher area. These have wavelengths of 1-4 m and heights of about 0.5 m.

Clay drapes that are deposited from suspension during falling water are common in the scour pits, and late ebb-stage, linguoid and parallel-crested ripples often occur on their stoss side and crest. Two box cores from between these dunes show that the preserved cross beds are flood-oriented and occur in sets 2-16 cm thick. Both planar and trough cross bedding is present. Mud drapes and silty mud balls up to 6 cm in diameter that were eroded from the channel bank are preserved. Significant changes in the structures may occur between spring and neap tides, but these have not been documented thoroughly.

Still higher on the bar, the dunes pass shoreward first into linguoid ripples and then into straight-crested ripples. The sands on this part of the bar are very fine-grained, and fluid mud drapes are present in low areas at low water. A box core from an area of straight-crested ripples on the highest part of the bar showed a set of ebb-oriented trough cross-bedding 8 cm thick that passes up into topset laminae that dip gently both down- and upriver. A box core from a linguoid rippled area at the upriver end of the bar showed ebb-dominant climbing ripples in cosets up to 6 cm thick. Clay drapes and organic debris were abundant. In the next bar downriver, dunes also pass laterally into linguoid and straight-crested ripples higher on the bar (Dickinson, 1991). The cross-bedding here is ebb-dominant, but alternates with parallel planar lamination that may have formed during upper-flow-regime conditions during flood tides.


Upriver Bars

The main accumulation of coarse-grained sediment in the inner estuary is found in the short reach immediately upriver from the Port Williams bridge (Fig. 6). This bar, which is composed of medium to very coarse-grained sand, is attached to the north bank at its higher, upriver end and separated from the north bank by a flood barb at its downriver end. This bar is best reached by climbing down the bank from the north dyke at its upriver high point. The medium dunes which cover this bar have wavelengths of about 4 m and heights of up to 40 cm. The concentration of coarse sand in this reach suggests that ebb and flood flow may be in balance here (i.e., ebb flow predominates upstream and flood ! flow predominates downstream from this reach). A box core from the upriver end of this bar showed alternating ebb and flood sets 3-19 cm thick. In contrast with the dunes on the Port Williams bar, silty mud drapes were preserved here.



The Upstream bar at the Port Williams Bridge

In the next reach upriver from this area of coarse sediment, a large bar of moderately well sorted, fine sand with an ebb barb is attached to the north bank of the river. Late ebb-stage ripples are developed on its surface. A box core from this bar showed mainly horizontal plane lamination, interrupted by a ripple laminated coset 10 cm thick. Ripple foresets 0.5-2 cm high dip alternately up-and downriver, but an ebb-orientation predominates. The prevalence of plane lamination indicates that upper plane bed conditions are common here. Bank-attached bars situated downriver of Port Williams also contain moderately well-sorted fine-grained sand and upper-flow-regime parallel lamination, but are flood dominated.



Tidal meanders upstream of the Port Williams Bridge

Stop 5: Kingsport Wharf (Fig. 1; Stop 5)

To the north you can see the eroding Triassic siltstone cliffs that provide the majority of sediment to the Cornwallis River and Avon River estuaries (Fig.7). The cliffs here are capped by glacial till and glacio-fluvial outwash. The influence of both waves and tidal currents is evident in the sequence of intertidal sediments. The high water mark is occupied by a wave-formed sandy beach which consists of material winnowed from the glacial sediments. Seawards, the sand gives way to either a bare, rock-cut platform (diagnostic of limited sediment supply), or a veneer of mudflat and occasional salt marsh. The mudflat grades seawards into sandy silt and thereafter into exposed bedrock. Mobile sand is found only in the base of tidal channels below the low water level. The bedrock exposed at low water shows a distinct bevel that is likely the result of wave erosion. The intertidal sediments in this region have a low preservation potential. The sand is slowly moved southwards around the government wharf and over the transgressed Acadian salt marsh, producing Oak Point (Fig. 7).

To the south of the government wharf a much thicker sedimentary succession is developed because of the sheltering provided by the wharf. A wave-formed beach is still present at the high water level, but this gives way outward to a well developed salt marsh and mudflat. The level of the salt marsh is artificially low due to the sheltering effects of the wharf.

Stop 6: Delhaven Wharf (Fig. 1; Stop 6)

A short stop at the Delhaven Wharf may provide us with a good opportunity for a photo depicting the full tidal range in the Minas Basin. Boats at this and other localities in the area have to use unique mooring techniques to compensate for the extreme tidal range. This is a good example of





Figure 7. Sketch map of Kingsport at the seaward limit of the Cornwallis estuary. Saltmarsh and Mudflats of the outer estuary give way between Oak Point and Kingsport wharf to a sediment-starved, wave-cut platform with a sand beach at the high tide level. This change reflects a seaward increase in the influence of wave action, and a change from tide dominated to mixed energy sedimentation.

one of the small fishing communities on the Minas Basin, the boats that dock here are inshore craft that exploit a wide variety of species (most commonly lobster) over the course of the year.



Fishing Boats at low tide at the Delhaven Wharf

Stop 7: The Lookoff (Figure 1; Stop 7)

The Lookoff affords a splendid view of Windsor Bay and the Cornwallis River-Avon River estuarine system; to the south, and of the eastern end of Annapolis Valley in which these estuaries are partially situated. The Annapolis Valley is located on the south limb of the plunging syncline of Triassic basalt that forms Cape Split, Cape Blomidon and the North Mountain. It is this feature that separates the wave dominated, high-energy outer Bay of Fundy from Minas Basin. On the seaward side, waves reaching 4 m in significant height have created an intertidal rock platform devoid of sediments. These waves in association with strong tidal currents are reworking the glacio-marine sequence which underlies much of the main Bay of Fundy (Fader *et al.*, 1977). The sand fraction has been moulded into large sand waves and mega ripples that are migrating northeastwards into Scott's Bay which occupies the centre of the syncline. The fines are swept through Minas Passage and into the Minas Basin where they accumulate in the inner reaches of the estuaries and embayments along the coastline.

The tidal amplification, and the associated increase in tidal current speeds, lead to a migration of the depocentre from the centre of the basin to the margins, with reworking of material in deeper water. Present-day bottom sediments of Windsor Bay show a progressive decrease in grain size landward from cobbles and boulders of the subtidal zone off Cape Blomidon, through gravelly sand to sand off Kingsport (Stop 5), and to silt off Evangeline Beach (Stop 3). Clay predominates still further up the Cornwallis River, beneath the turbidity maximum.

REFERENCES

- Amos, C.L. 1978: The post glacial evolution of the Minas Basin, N.S.: a sedimentological interpretation; *Journal of Sedimentary Petrology*, v.48, p.965-982.
- Bleakney, J.S. and Davis, D. 1983: Discovery of an undisturbed bed of 3800 year old oysters (*Crassostrea virginica*) in Minas Basin, Nova Scotia. *Proceedings of the Nova Scotia Institute of Science*, v.33, p.1-6.

- Daborn, G. (ed.) 1991: Littoral Investigation of Sediment Properties - LISP 89 Final Report; Acadia Centre for Estuarine Research, Publication 17, 239 p.
- Daborn, G.R. and Pennachetti, C. 1979: Physical oceanographic and sedimentological studies in the Southern Bight of Minas Basin. Proceedings of the Nova Scotia Institute of Science, v. 29, p.315-333.
- Dalrymple, R.W., Amos, C.L. and McCann, S.B. 1982: Beach and Nearshore Depositional Environments of the Bay of Fundy and Southern Gulf of St. Lawrence; International Association of Sedimentologists, 11th Congress, Hamilton, Guidebook Excursion 6A, 116 p.
- Dickinson, A. 1991: Sedimentary structures of the Cornwallis River; B.Sc. thesis, Acadia University, Wolfville, Nova Scotia. 124 p.
- Fader, G.B., King, L.H. and McLean, B. 1977: Surficial Geology of the Eastern Gulf of Maine and Bay of Fundy; Geological Survey of Canada, Marine Science Paper 76-17, 23 p.
- Ferguson, S. A. 1983: Geological map of the Hantsport area; Nova Scotia Department of Mines and Energy, Map 83-1 (1:25 000 scale)
- Goldthwaite, J.W. 1924: Physiography of Nova Scotia; Geological Survey of Canada, Memoir 140, 179 p.
- Harrison, W. and Lyon, C.J. 1963: Sea Level and crustal movements along the New England - Acadian shore, 4500-3000 B.P.O.; Journal of Geology, v.71, p.96-108.
- Hubert, J.F. and Mertz, K.A. 1980: Eolian dune field of Late Triassic age, Fundy Basin, Nova Scotia. Geology, v.8, p.516-519.
- Klein, G. deV. 1962: Triassic sedimentation, Maritime Provinces, Canada ; Bulletin of the Geological Society of America, v.73, p. 1127-1146.
- McDow, N.C. 1991: The stratigraphy and sedimentology of the Guzzle and Boot Island, Minas Basin, Bay of Fundy; B.Sc. thesis, Acadia University, Wolfville, Nova Scotia, 120 p.
- Reineck, H.-E. and Singh, I.B. 1980: Depositional Sedimentary Environments (2nd edition); Springer-Verlag, New York, 549 p.
- Schubel, J.R. (ed) 1971: The Estuarine Environment, Estuaries and Estuarine Sedimentation; Wye Institute, Short Course Lecture Notes, 30-31, October 1971, 349 p.