

**DEFORMATION AND METAMORPHISM IN THE CONTACT
AUREOLE OF THE SOUTH MOUNTAIN BATHOLITH, BEDFORD AREA**

Paul D. Gray

Submitted in Partial Fulfillment of the requirements
for the Degree of Bachelor of Science, Honors
Department of Earth Sciences
Dalhousie University, Halifax, Nova Scotia
March 1996



Dalhousie University

Department of Earth Sciences

Halifax, Nova Scotia

Canada B3H 3J5

(902) 494-2358

FAX (902) 494-6889

DATE April 9 1996

AUTHOR Paul D. Gray

TITLE Deformation and Metamorphism in the Contact
Aureole of the South Mountain Batholith,
Bedford Area

Degree B.Sc Convocation May Year 1997

Permission is herewith granted to Dalhousie University to circulate and to have copied for non-commercial purposes, at its discretion, the above title upon the request of individuals or institutions.

THE AUTHOR RESERVES OTHER PUBLICATION RIGHTS, AND NEITHER THE THESIS NOR EXTENSIVE EXTRACTS FROM IT MAY BE PRINTED OR OTHERWISE REPRODUCED WITHOUT THE AUTHOR'S WRITTEN PERMISSION.

THE AUTHOR ATTESTS THAT PERMISSION HAS BEEN OBTAINED FOR THE USE OF ANY COPYRIGHTED MATERIAL APPEARING IN THIS THESIS (OTHER THAN BRIEF EXCERPTS REQUIRING ONLY PROPER ACKNOWLEDGEMENT IN SCHOLARLY WRITING) AND THAT ALL SUCH USE IS CLEARLY ACKNOWLEDGED.

Distribution License

DalSpace requires agreement to this non-exclusive distribution license before your item can appear on DalSpace.

NON-EXCLUSIVE DISTRIBUTION LICENSE

You (the author(s) or copyright owner) grant to Dalhousie University the non-exclusive right to reproduce and distribute your submission worldwide in any medium.

You agree that Dalhousie University may, without changing the content, reformat the submission for the purpose of preservation.

You also agree that Dalhousie University may keep more than one copy of this submission for purposes of security, back-up and preservation.

You agree that the submission is your original work, and that you have the right to grant the rights contained in this license. You also agree that your submission does not, to the best of your knowledge, infringe upon anyone's copyright.

If the submission contains material for which you do not hold copyright, you agree that you have obtained the unrestricted permission of the copyright owner to grant Dalhousie University the rights required by this license, and that such third-party owned material is clearly identified and acknowledged within the text or content of the submission.

If the submission is based upon work that has been sponsored or supported by an agency or organization other than Dalhousie University, you assert that you have fulfilled any right of review or other obligations required by such contract or agreement.

Dalhousie University will clearly identify your name(s) as the author(s) or owner(s) of the submission, and will not make any alteration to the content of the files that you have submitted.

If you have questions regarding this license please contact the repository manager at dalspace@dal.ca.

Grant the distribution license by signing and dating below.

Name of signatory

Date

ABSTRACT

This study documents structures related to Halifax Pluton emplacement, in the Bedford area. Intrusion of the Devonian South Mountain Batholith caused both metamorphism and deformation of the Cambrian-Ordovician Meguma Group. Most previous structural interpretations for the area have attributed local structural features to regional deformation events. Field and petrographic analysis indicates that a strain and thermal aureole exists in a zone surrounding the Halifax Pluton. Mapping revealed several structures hitherto locally unrecognized, including a schistose lineated fabric (LS_2) and other minor contact-parallel features. Microscopic analysis of these rocks defined an aureole at least 1.5 km wide, divided into an outer biotite zone, a medial cordierite zone, and an inner K-feldspar zone. A cross-section along the axial trace of the Bedford Syncline indicates a variation in style of deformation and strain with proximity to the granite contact. An increase in dip of bedding towards the contact is interpreted as a result of flattening perpendicular to the granite boundary, probably with a high ratio of pure shear to simple shear. The same cross-section also offered insight on the depth to detachment of both the transverse anticline and the larger Magazine Hill Basin. Metamorphic minerals define the deformation structures (e.g. LS_2), which suggests deformation and metamorphism were both imposed by the rising, and possibly ballooning granite pluton. Therefore this study proposes that metamorphism and significant deformation of the Goldenville Formation in the study area are related spatially and temporally to the emplacement of the Halifax Pluton.

Keywords: Halifax Pluton, Goldenville Formation, Contact Metamorphism, Deformation, Emplacement, Aureole, Foliation

ACKNOWLEDGEMENTS

I would like to thank my advisors Dr. Nicholas Culshaw and Dr. Rebecca Jamieson for their continual, informative assistance. I also thank Dr. Martin Gibling for his grammatical and organizational help. Most importantly without my fathers advice and enthusiasm this thesis would not have been possible. Lastly, my mother and Robbie Shepard gave constant necessary support, which as well made this work possible.

TABLE OF CONTENTS

	Page
Abstract	i
Acknowledgements	ii
Table of Contents	iii
Table of Figures	v
Chapter 1 Introduction	1
1.1 Introduction	1
1.2 Pluton Emplacement	2
1.3 Emplacement Effects	4
1.4 Scope and Purpose	7
1.5 Organization	9
Chapter 2 Regional Geology	10
2.1 Introduction	10
2.2 Meguma Group	12
2.3 Acadian Deformation of the Meguma Group	13
2.4 Batholith Emplacement	14
2.5 Post-Granite Deformation	15
2.6 Summary	16
Chapter 3 Field Relations	17
3.1 Introduction	17
3.2 Thermal and Strain Aureoles	17
3.2.1 Metamorphic Thermal Aureole	17
3.2.2 Strain Aureole	20
3.3 Major Structures in the Kearney Lake Area	21
3.4 Predominant Minor Structures	22
3.5 Other Minor Structures	25
3.5.1 Boudins	25
3.5.2 Pegmatite	27
3.5.3 Extensional Planar Features	27
3.6 Eastern Pluton Edge	29
3.7 Summary	30

Chapter 4 Microscopic Structures and Metamorphism	31
4.1 Introduction	31
4.2 Metamorphic Aureole Zonation	33
4.2.1 Biotite Zone	33
4.2.2 Cordierite Zone	34
4.2.3 K-feldspar Zone	39
4.3 Similar Contact Aureoles	41
4.4 Timing of Metamorphism and Deformation	48
4.5 Conclusions	49
Chapter 5 Discussion and Conclusions	52
5.1 Introduction	52
5.2 Cross-Section	53
5.3 Strain Path at the Granite Contact	55
5.4 Field and Microscopic Interpretations	58
5.5 Halifax Pluton Emplacement	58
5.6 Conclusions	59
References	61
Appendix A: Table of Field Data Compilation (Fieldlog Database)	A1
Appendix B: LS₂ Zone Measurements	B1
Appendix C: Table of Petrographic Descriptions	C1

TABLE OF FIGURES

	Page
Chapter 1	
Figure 1.1 Pluton Emplacement Mechanisms	3
Figure 1.2 Foliation Patterns Around Temporally Distinct Plutons	5
Figure 1.3 Location Map	8
Chapter 2	
Figure 2.1 Regional Geological Map	11
Chapter 3	
Figure 3.1 Simplified Structural Geological Map of Study Area	18
Figure 3.2 Major Geological Structures in Study Area	19
Figure 3.3a Photo Illustrating LS ₂ Fabric	23
Figure 3.3b Block Diagram Illustrating LS ₂ Fabric	23
Figure 3.4 Schistose S ₂	24
Figure 3.5 Calc-silicate Boudins with Reaction Rims	26
Figure 3.6 Cross-Cutting Pegmatite	28
Chapter 4	
Figure 4.1 Sample Sites and Metamorphic Isograds	32
Figure 4.2 Typical Biotite Zone Sample	35
Figure 4.3 Randomly Oriented Cordierite and Biotite Porphyroblasts	37
Figure 4.4 Intensified Foliation at Porphyroblast Edges	38
Figure 4.5 Grain Coarsening in K-feldspar Zone	40
Figure 4.6a Wavy Biotite Bands and Intergranular Cordierite	42
Figure 4.6b Aligned Biotite and Intergranular Cordierite	43
Figure 4.7a Ring-like Structures	44
Figure 4.7b Biotite Grains Defining Rings	45
Figure 4.8 Myrmekite Alteration	46
Figure 4.9 Summary of Mineral Assemblages	50
Figure 4.10 Phase Diagram	51
Chapter 5	
Figure 5.1 Cross-Section	54
Figure 5.2 Schematic representations of Calculation Results	56

CHAPTER 1 INTRODUCTION

1.1 Introduction

The South Mountain Batholith (SMB) forms one of the largest intrusive granitoid complexes in the Appalachian Orogen (Abbott, 1989). Emplacement of the South Mountain Batholith resulted in metamorphism and deformation of the pre-existing country rocks of paleo-Nova Scotia, the Meguma Group. This thesis concerns the Halifax Pluton member of the South Mountain Batholith, and its effect on the Meguma Group during emplacement.

The emplacement of the Halifax Pluton formed a metamorphic contact aureole adjacent to the pluton. In the Bedford area, this aureole contains a number of features never previously described, including metamorphic foliation, stretching lineation, and boudins. This study documents these features from structural, metamorphic, mesoscopic, and microscopic standpoints. A synthesis of these data allows new interpretations on the deformation, displacement, and metamorphism at the pluton boundary.

1.2 Pluton Emplacement

Pluton emplacement involves a body of magma upwelling through the crust and associated wall rock deformation and metamorphism. Mechanisms of magma rise have been the subject of vigorous debate since the 1940's. In recent years, the focus has changed to the connection between plutonism and regional deformation. Terms like "passive" (no forceful injection) and "active" (forceful injection) emplacement have been adopted without much thought about the mechanics behind such nomenclature (Paterson and Fowler, 1993).

The structural effects of pluton emplacement can change character with depth, even within a single pluton. Near surface, epithermal plutons are characterized by brittle deformation of wall rocks, moderate depth, mesothermal plutons are associated with transitional (brittle to ductile) deformation, whereas deep hypothermal interfaces show ductile deformation (Woodcock and Underhill, 1987; Paterson et al., 1992).

Stoping, cauldron subsidence, ballooning, diapirism, and doming are all possible mechanisms of pluton emplacement, and all have distinctive deformation patterns (Paterson et al., 1991b) (Fig. 1.1). Rise of magma is driven by density contrasts between the melt and surrounding rocks. The melt is more buoyant than the surrounding rocks, and this contrast drives the magma up (Paterson and Vernon, 1995). Rising ceases when the magma achieves a position of dynamic equilibrium in the crust. The crustal level where equilibrium is acquired is a complex function of the density, competency, and




Figure 1.1 Pluton emplacement mechanisms. 1. Stopping. 2. Roof doming, with associated block faults. 3. Return flow and ductile wall rock alteration. 4. Anatexis and assimilation. 5. Lateral wall rock displacement. (From Paterson et al., 1991).

rheology of both the magma and the surrounding country rock. Therefore emplacement is controlled by a combination of these factors (Paterson and Fowler, 1993).

1.3 Emplacement Effects

Wall rocks and magma interact as a coupled, open, and linked system, with chemical and physical, e.g. material, exchange across the granite-wall rock boundaries. The emplacement of granite is affected by the wall rocks, and vice versa (Paterson et al., 1991a). In ideal conditions, a chronology of events related to pluton emplacement can be deduced from features recorded in metamorphic aureoles.

The contact temperatures country rocks are subjected to during pluton emplacement can be intense enough to cause melting. This melting results in a metamorphic contact aureole, which while not regionally important, can be locally extensive (Paterson et al., 1993). Contact aureoles are important because they may preserve a recordable gradient of the interaction of magma with surrounding rocks. Metamorphosed wall rocks, if properly interpreted, offer an unrivalled record of conditions during emplacement (Paterson et al., 1991b).

Perhaps the most pertinent information that aureoles can convey concerns the timing of pluton emplacement. Relative timing of regional events (folding, faulting, or cleavage development), with respect to pluton emplacement, can be deduced from textural relationships in contact aureoles (Fig. 1.2). A pluton emplaced after regional deformation would overprint regional fabrics within the aureole. On the other hand, a pluton emplaced before regional deformation may affect subsequent




Figure 1.2 Hypothetical foliation patterns around pre- (a,d), syn- (b,e), and post-tectonic (c,f) plutons. (From Paterson et al., 1991).

deformation by deflecting it, or obviously be affected by it. Therefore, suitable aureoles may indicate relative age determinations like pre-, syn-, or post-tectonic emplacement (Bateman, 1985; Guglielemo, 1994; Paterson et al., 1991b,1993).

The interpretative value of metamorphic contact aureoles is well understood. Contact aureoles offer more than just interesting "natural laboratories" for unravelling metamorphic reactions. The kinematics and mechanisms behind the development of structures, such as folds and foliations, may also be inferred. Well documented contact aureoles are renowned for their emplacement-related metamorphic foliations, faults, folds, lineations, and high-grade metamorphism of adjacent rocks (Bateman, 1985; Woodcock and Underhill, 1987; Fyson, 1980; Paterson et al., 1991a). Contact aureoles offer an indication of gradation in metamorphism over a relatively small area. Relationships between structures such as fabrics, folds, and faults can be studied in these small, well constrained areas.

Deformational fabrics in aureoles offer unique insights into emplacement-related strain paths and histories. Pluton emplacement is a "destructive" event, resulting in high-grade metamorphism and possibly complete annealing of pre-existing fabrics in country rocks.

Halifax Pluton emplacement appears to have been directly responsible for contact-parallel foliation, steeply plunging lineations, the transverse Kearney Lake Anticline, and possibly the Magazine Hill Basin of Burns (1995).

1.4 Scope and Purpose

This study documents and describes deformation of the Cambrian-Ordovician Meguma Group at the northeast margin of the Devonian South Mountain Batholith, Halifax Pluton. Both structural and metamorphic aspects of pluton-related deformation are addressed. Particular attention is paid to contact metamorphism and related foliation development within the pluton-adjacent rocks.

Mesoscopic and microscopic analysis of mineralogy and mineral assemblages of the metamorphosed Goldenville Formation are presented.

The study area lies west of the Bedford Basin, north of Halifax, Nova Scotia. Most field work was done in the Kearney Lake area, just west of exit 2, off Highway 102 (Fig. 1.3). This area was chosen because it allowed a follow-up to Burns (1995) study.

Numerous workers have described the geology of the general area (Faribault, 1908; Horne et al., 1992; Clarke and Chatterjee, 1988). Yet the contact zone has several distinct structures previously undocumented. A series of recent road-cuts has allowed investigation of the contact zone. The transverse anticline located close and parallel to the Halifax Pluton contact was noted in previous studies (Faribault, 1908; Clarke and Tate, 1993), as was the more distant Magazine Hill Basin described by Burns (1995). Burns proposed a possible emplacement related derivation of the transverse folds and the Magazine Hill Basin.

This thesis examines effects of the emplacement of the Halifax Pluton from a number of data sources. These data sources include field mapping aided by Global Positioning equipment (Trimble Pathfinder+ with differential corrections, allowing < 5 meter accuracy), use of thin sections prepared




Figure 1.3 Location map of study area. Accompanying legend for use with all maps to follow, unless otherwise indicated.

from oriented samples collected within the aureole, and newly acquired measurements of bedding (S_0), slaty cleavage (S_1), metamorphic foliation (S_2), and stretching lineations (L_2). A compilation of these newly acquired data, plus previous measurements (Appendix A), allows a structural and metamorphic synthesis of the effects of pluton emplacement.

1.5 Organization

Chapter 2 deals with the regional geological setting, as well as local geological evolution. Chapter 3 describes the macroscopic structures and features within the contact hornfels of the Goldenville Formation, including the regional distribution, geometry, nature, and morphology of metamorphic fabrics. Chapter 4 describes the metamorphic nature of the contact aureole from a microscopic standpoint, with thin section descriptions. Chapter 5 presents new structural interpretations from collected data and computer aided calculations, and concludes this thesis.

CHAPTER 2 REGIONAL AND LOCAL GEOLOGY

2.1 Introduction

Atlantic Canada, and the Canadian Appalachians, contain several distinct terranes, including the Avalon, Gander, and Humber Terranes. Within Nova Scotia, the Avalon and Meguma Terranes are separated by the extensive strike-slip, Cobequid-Chedabucto Fault System (Fig. 2.1). The Meguma Terrane is dominated by two rock types: Cambrian-Ordovician Meguma Group metasedimentary rocks and Devonian granitoid bodies. This thesis is concerned with both the granitoid bodies (the South Mountain Batholith) and the Meguma Group.

The Meguma Terrane was metamorphosed and deformed during the Mid-Late Devonian Acadian Orogeny and metamorphosed again during subsequent plutonism. Timing and relationship of orogeny and plutonism is unclear, however evidence indicates that deformation continued in an episodic fashion after pluton emplacement (Horne and Culshaw, 1993). The orogeny may have played a pivotal role in controlling emplacement of the granite. These controlling factors will be addressed in following chapters.




Figure 2.1 Regional geological map of Nova Scotia (Modified from Keppie, 1987)

2.2 Meguma Group

The Meguma Group contains the oldest rocks within the Meguma Terrane: Cambrian-Ordovician metapelites and metapsammites of the Halifax and Goldenville Formations, respectively. Together they form the largest single mass of sedimentary rock in the Atlantic Provinces. The older Goldenville Formation is a psammitic flysch-like deposit with an approximate stratigraphic thickness of six kilometers. The younger Halifax Formation is a pelitic shaly-flysch with a similar six kilometer thickness (Schenk, 1981). Together with the overlying Silurian-Devonian sedimentary and volcanic rocks, Schenk (1991) has estimated a total stratigraphic thickness of 20 km.

The Meguma Group is overlain by metaquartzite and other lithologies of the White Rock Formation, the shaly to silty Kentville Formation, the volcanoclastic New Canaan Formation, and the shaly to silty Torbrook Formation. These formations have depositional ages estimated from Ordovician-Silurian to Devonian (Emsian) (Schenk, 1981). The South Mountain Batholith intruded the Meguma Terrane between the Emsian and the Tournaisian, at approximately 370-395 Ma (MacDonald et al., 1992; Clarke et al., 1993). The Horton Group (Late Devonian to Tournaisian age) is a variable clastic terrestrial sediment, which non-conformably overlies the South Mountain Batholith.

Meguma Group deposition in a deep-sea to near-shelf gradational fan complex is one of the more widely accepted depositional theories (Schenk, 1981). In this setting, the Halifax Formation most likely represents a deep-water, anoxic depositional environment, whereas the

Goldenville Formation represents a more sandy, closer-to-shore environment. Schenk (1991) has further argued for deposition on the continental margin of Gondwana.

2.3 Acadian Deformation of the Meguma Group

The Meguma Group and overlying Silurian-Devonian White Rock, New Canaan, Kentville, and Torbrook Formations were deformed and metamorphosed during the Acadian Orogeny (Williams, 1979; Keppie and Dallmeyer, 1987; Horne et al., 1992). This deformation resulted in the development of straight limbed, open to tight, northeast-southwest trending regional folds, with associated axial planar cleavage (Faribault, 1908; Henderson et al., 1986; Horne and Culshaw, 1993). Evidence suggests that this deformation occurred ca. 390 Ma, prior to the intrusion of the peraluminous granitoid bodies at ca. 375 Ma (Clarke et al., 1993).

Theories abound as to the cause and timing of deformation of the Meguma Group (Lajtai and Stringer, 1981; Keppie, 1982; Kontak et al., 1989). If the Meguma is indeed an Acadian accreted terrane, as widely accepted, deformation may have been directly associated with docking during the Acadian orogeny. Later deformation, which undoubtedly occurred (Horne and Culshaw, 1993), would entail reactivated regional folds and enhanced folding and cleavage development (pers. comm., N Culshaw, 1995).

2.4 Batholith Emplacement

About half the area of the southern peninsula of Nova Scotia is underlain by the South Mountain Batholith (Fig. 2.1), at over 7300 km² the largest exposed peraluminous granitoid body of the Appalachian Orogen (MacDonald et al., 1992). It is about 180 km long and 50 km wide. Abbott (1989), using geophysical information, interpreted the average thickness of the South Mountain Batholith as 5 km, with a maximum thickness of 25 km near the New Ross area.

MacDonald et al. (1992) subdivided the South Mountain Batholith into thirteen plutonic members, associated with two stages of emplacement. Early plutons (Stage 1), comprise granodiorite and monzogranite. Later plutons (Stage 2), comprise mostly monzogranite, leucomonzogranite, and leucogranite. The Halifax pluton, the focus of this study, is a stage 2 pluton.

The South Mountain Batholith intruded the Meguma Group later than the regional deformation event (Horne et al., 1992; Clarke et al., 1993), however, a connection between the two events is reasonable. Two possible causes of plutonism are: increased crustal thickness, and subduction-related lithosphere removal. After accretion, the increased crustal thickness may have caused melting at depth, with subsequent magma upwelling. This model, however, may not be feasible, due to the limited thickness of the Meguma (some 20 km), compared with crustal depth. With such constraints, the increase in crustal thickness would not be enough to cause melting at depth (pers. comm., N. Culshaw, 1995). Clarke and Tate (1993) postulated that the subducting North African plate caused lithospheric plate removal under paleo-Nova Scotia. This removal

allowed upwelling of the hot asthenosphere, allowing crustal melting. The ensuing crustal melt combined with mantle melt, forming the South Mountain Batholith and small ultramafic to mafic intrusions.

Studies on the Liscomb Complex by Clarke et al. (1993) have shown that the granite source may be sub-Meguma basement rock. Crustal melting and crustal assimilation seem to be involved with granite generation at depth.

Horne et al. (1992) have produced some evidence of structural controls on South Mountain Batholith emplacement. This evidence includes the two main pluton boundary orientations, northeast and northwest. Pre-existing faults probably influenced stage 2 pluton emplacement. Further, the regional similarity of joint trends in second stage plutons possibly indicates some degree of syntectonic emplacement. There may have been a continued sense of stress applied to the region during and after this second stage of emplacement.

2.5 Post-Granite Deformation

Post-South Mountain Batholith deformation is evident throughout the Meguma Terrane. Reactivation of existing faults and folds, complete with intensification of cleavage, is the result of this later deformation (Horne and Culshaw, 1993). Later deformation could explain the "wrap around" cleavage seen along the southeast margin of the South Mountain Batholith. Deformation may have occurred at such low temperatures (lower greenschist facies) that the granite was only slightly (non-visibly) altered.

Horne and Culshaw (1993) argue convincingly for post-granite deformation. Their evidence includes flexural slip shear fractures within the Halifax Formation that truncate cordierite porphyroblasts associated with the thermal aureole of the Halifax pluton in Halifax. Some deformation certainly took place post-South Mountain Batholith emplacement, but how long after is unclear.

2.6 Summary

Regional folds and axial planar cleavage were produced in the Meguma Terrane during the Devonian Acadian Orogeny. Relatively shortly thereafter, plutonism occurred, forming the South Mountain Batholith. Deformation continued after, and possibly during South Mountain Batholith emplacement, controlling its emplacement. Evidence of direct deformational effects of pluton emplacement are seen only in the immediate vicinity of the pluton, and no extensive regional structures resulting from granite emplacement have been documented. Structures resulting from deformation are described in the next two chapters.

CHAPTER 3 FIELD RELATIONS

3.1 Introduction

This chapter describes the mesoscopic and macroscopic features at the northeast end of Halifax Pluton, within and close to the Halifax Pluton metamorphic aureole. There are several major and minor structures within the aureole; none however are evident in the granite. The major structures are: the northeast-southwest (trend 230°) regional folds (Waverley Anticline and Bedford Syncline), the transverse (contact-parallel) Kearney Lake Anticline (trend 120°), and the Magazine Hill Basin described by Burns (1995) (Fig. 3.1 and 3.2). Minor structures include: axial planar cleavage of the regional folds (S_1), metamorphic foliation (S_2), mineral lineations (L_2) (together forming a composite fabric, LS_2). The major fold-structures illustrated in Figure 3.1 are defined by bedding (S_0) trends measured by the author, and those compiled from Faribault (1908) and Burns (1995).

3.2 Thermal and Strain Aureoles

3.2.1 *Metamorphic Thermal Aureole*

A metamorphic contact aureole is defined as a zone surrounding an igneous intrusion, characterized by contact metamorphism (Yardley, 1989). The metamorphic thermal aureole of the Halifax Pluton contains contact metamorphic features which grade away from the pluton into




Figure 3.1 Major geological structures and simplified structural geology of study area and vicinity.




Figure 3.2 Geology of the Kearney Lake area. Interpreted Magazine Hill Basin of Burns (1995) is shaded. LS₂ schist zone is stippled. A-F line represents cross-section on Figure 5.1. Inset stereonet: shear band-like planar features (thin great circles) and S₀ (thick great circle).

regional metamorphic features. The change is reflected in textural and mineralogical distinctions, e.g. appearance of cordierite and associated grain coarsening. The thermal aureole extends about 1.5 km from the granite contact. Chapter 4 will cover this topic.

3.2.2 *Strain Aureole*

The strain aureole of the Halifax Pluton is defined here as the zone containing fabrics directly related to pluton emplacement. The schist with planar-linear fabric (LS_2), the transverse Kearney Lake Anticline, and the Magazine Hill Basin, are all interpreted as emplacement related structures, because of their anomalous nature compared to regional structures of the Meguma Group (Fig. 2.1). Interestingly, the strain aureole extends farther than the thermal aureole, approximately six kilometers.

The transect on Figure 3.2 illustrates the strain aureole, the variation of rock types within it, and the aforementioned major structures. The point labelled (A) lies within the unfoliated, massive monzogranite. The contact between the granite and hornfels is indicated by point (B). The rocks here contain a well developed, penetrative, planar-linear schistose fabric, LS_2 . LS_2 decreases in intensity with distance from the granite, and is less penetratively developed between C and D. At approximately point D, LS_2 is no longer mesoscopically discernible, and axial planar cleavage (S_1 , defined by phyllosilicates) appears in slates. This trend continues farther out, with S_1 intensifying. Point E represents the beginning of the Magazine Hill Basin of Burns (1995).

3.3 Major Structures at the Northeast End of Halifax Pluton

Figures 3.1 and 3.2 illustrate the major structures in this study: the Waverley Anticline, the Bedford Syncline, the Magazine Hill Basin of Burns (1995), and the transverse Kearney Lake Anticline. Field mapping from this study has relocated granite contact placement slightly from the provincial map.

Bedding which defines these major structures, measured in the Goldenville Formation is difficult to recognize, particularly in the rather massive, recrystallized hornfels of the contact aureole. If present, rare metamorphosed shaly interbeds however, provide reliable measurements.

The transverse Kearney Lake Anticline trends parallel to the granite boundary (approximately 120°), and normal to the regional folds. This fold has an approximate length of >4.3 km and width of 1.6 km. It is an upright fold, with an approximate interlimb angle of 130° . The regional Bedford Syncline in comparison trends 270° , is periclinal, upright, has an approximate interlimb angle of 150° , a cylindrical axial trace for tens of kilometers, and is about 3.5 kilometers wide (Faribault, 1908).

Previous studies in the area (Faribault, 1908; MacDonald et al., 1992; Burns, 1995), indicated the presence of a transverse rim syncline associated with the Kearney Lake Anticline. The author has inspected the area, and was unable to recognize bedding in outcrops in the critical area of the reported syncline. Faribault's measurements have however been confirmed with astounding accuracy at almost every location. The syncline may be present, but it will require



further careful field work to confirm or disprove its existence. Hence, with all due respect, this study proposes then that the rim syncline is not reliably defined.

3.4 Predominant Minor Structures

Predominant minor structures in the strain aureole are; the metamorphic schistosity (S_2 component of the LS_2 fabric) and the steeply plunging mineral lineation (L_2). L_2 lies in the S_2 plane, and together with S_2 defines the LS_2 schist (Fig. 3.3a). LS_2 contains no asymmetric fabric elements, e.g. rotated feldspars, or C-S fabrics. The variably developed S_2 parallels the trend of the granite boundary, striking approximately 120° . The width of the LS_2 schist band ranges up to 250 m, and is developed over the entire length of the contact in the study area, some 10 km (Fig. 3.2). Most common orientations of S_0 and S_2 are $100^\circ/80^\circ$ and $120^\circ/85^\circ$, respectively (Appendix B). In a very narrow zone within this schist band, close to the contact, S_0 and S_2 are sub-parallel to parallel. Beyond approximately 250 m from the contact, schistosity is indistinct.

A shift in strike accompanies the decrease in LS_2 intensity. Within the LS_2 zone, foliation is essentially parallel to the contact. Roughly 300 m from the contact, the strike of S_2 (230°) resembles that of regional fold limbs (Fig. 3.2). Intermediate trends are measured between 300 and 700 m from the contact. No evidence of over-printing between S_1 and S_2 was observed.

Hand samples of LS_2 display compositional banding defined by alternating leucocratic (quartz-feldspar-cordierite) and melanocratic (biotite) aggregates at a mm scale (Fig. 3.4). These leucocratic aggregates are penetrative, and have a flattened, blade-like shape. Individual grains



LEGEND

	- L_2 lineations
	- Boudinage (Calc-silicate Nodules)
	- Pegmatite
	- S_2 Foliation

Figure 3.3a) Photograph of LS_2 fabric from site 95-76. Steeply dipping lineation (L_2) parallel to pencil, metamorphic foliation (S_2), an S_2 parallel band containing L_2 , and boudins (stretched along S_2 and down L_2). Pencil is 14 cm. b) Idealized block diagram illustrating LS_2 fabric.




Figure 3.4 Weathered surface showing schistose S_2 (sample 95-15). Although foliation appears curved, weathered surface is curved, giving false impression of plane foliation. Red bar is 1 cm.

defining the aggregates are small, ~1 mm on average. Complete recrystallization is evident in these rocks, with weak grain elongation or alignment. Chapter 4 will investigate this aspect of these LS_2 fabrics.

Away from the contact the well developed LS_2 gives way to a non-penetrative fabric. These subtle fabrics are best recognized on weathered surfaces. In thin section, this fabric is not well defined because the scale of aggregates that define the fabric are greater than a low power field of view, 3-5 mm.

3.5 Other Minor Structures

Within the LS_2 schist zone, other minor structures occur, including boudins and quartz and feldspar (granite ?) filled extensional, shear-band-like structures.

3.5.1 Boudins

Decimetric boudins are developed from cross-cutting and concordant pegmatites and calc-silicate layers. The blade-like geometry of the boudins is illustrated in Figures 3.3a/b and 3.5. They display elongation within the S_2 plane, both parallel to the strike of S_2 and the plunge of L_2 . A flattened, blade-like shape is thus evident (K value between 1 and 0). This shape fabric indicates a similar strain geometry to LS_2 , albeit a larger scale.



Figure 3.5 Calc-silicate boudins, on a horizontal surface. Necks of calc-silicate join separate boudins, and slightly darker reaction rims surround the boudins. Diameter of lens cap 5 cm.

Calc-silicate layers that form many of the boudins, are common in some stratigraphic horizons of the Goldenville Formation. The shape of these boudins is comparable to the shape fabric that occurs in the LS₂ schist, and is consistent with the finite strain of these rocks. Close to the contact (100-150 m), foliation wraps around the calc-silicate nodules in an augen-like manner, suggesting ductile flow of the country rock. Reaction rims of slightly darker minerals, averaging <5 mm widths, form concentric zones on these nodules, making them stand out on weathered surfaces (Fig. 3.5).

3.5.2 Pegmatite

Two orientations of pegmatite exist in the LS₂ schist zone: concordant to LS₂ (boudinaged) approximate trend 115°, and cross-cutting (filling fractures perpendicular to the lineation) approximate trend 190° (Fig. 3.3b and 3.6). The presence of such kinematically early and late pegmatite indicates that granite was available throughout the period of deformation. This evidence indicates syn-plutonic deformation of the country rocks.

3.5.3 Extensional Planar features

Quartz and feldspar (?) filled planar features are common immediate to the contact. They displace L₂, and they cut and separate the schistosity at approximately 280° (Appendix B). These dip away from the granite and occupy a very narrow zone, 25-35 m, from the contact (Fig. 3.2).



Figure 3.6 Cross-cutting pegmatite on horizontal surface, cutting and separating S_2 and calc-silicate boudins. Diameter of lens cap 5 cm.

3.6 Eastern Pluton Edge

This study focused on the northeast end of the Halifax Pluton, and its contact aureole. Some mapping and sampling was done farther south, around the corner of the pluton, near Burnside Industrial Park (Fig. 3.3). S_2 and L_2 are present in this area, though much less well developed. S_2 orientations vary considerably from those at the northeast margin, with averages close to regional trends, 240° . Also, the granite-country rock contact is more irregular in detail, than at the northeast margin.

These distinctions from the northeast end of the pluton may be a indicative of specific emplacement-related processes. For example, if the pluton was expanding normal to regional compression, the northeast end would show signs of forceful injection, e.g. contact-parallel LS_2 . In the same scenario, sides parallel to regional compression would not display the same well developed features (Guglielmo, 1992). The study area is not extensive enough to answer these questions, however, documentation of these observations may prove important for further studies.

3.7 Summary

Major and minor structures occur within the strain aureole of the Halifax Pluton. Structures close to the contact (e.g. LS_2 and the Kearney Lake Anticline), trend parallel to the contact, unlike any other regional structures. The anomalous nature of these structures, the presence of syn-kinematic pegmatites, and the obvious relationship between metamorphic

recrystallization and LS_2 schistosity, leads to the interpretation that granite emplacement directly caused their development.

The shape of the finite strain ellipsoid, deduced from various features (Fig. 3.3a and b), reflects the integrated total strain accompanying the deformation. The strain shape, regardless of scale (LS_2 , boudins, or S_2), is a triaxial, flattened, blade-like ellipsoid, with its long axis near vertical and a component of extension parallel to the strike of the granite contact. Further, the lack of asymmetric fabric elements associated with LS_2 , suggests irrotational, pure flattening strain. A more quantitative view of this evidence is the topic of chapter 5.

CHAPTER 4 MICROSCOPIC STRUCTURES AND METAMORPHISM

4.1 Introduction

This chapter describes the microscopic structures and features within the metamorphic contact aureole of the Halifax Pluton. Petrographic descriptions are based on thin sections collected and prepared by the author (Appendix C). Sample sites, indicated on Figure 4.1, represent traverses across the aureole, from regional structures to the pluton. Data compiled from these traverses illustrate the variation in mineralogy and textures, embodied as changes in metamorphic grade within the Goldenville Formation adjacent to the Halifax Pluton.

A thermal contact aureole, extending approximately >1.5 kilometers from the pluton-Goldenville Formation contact is here defined on the basis of microscopic features. As is common in contact aureoles, contact metamorphic features grade directly into regional metamorphic features (Pattison and Tracy, 1991), hence an accurate outer limit of the aureole was not defined in this study area. The mesoscopic features used to define the width of the aureole (Chapt. 3), are augmented by microscopic evidence in the present chapter.

The thermal aureole of the Halifax Pluton exhibits several attributes typical of contact metamorphic aureoles. Kerrick (1991) defined contact aureoles in terms of processes that the country rocks were subjected to during granite intrusion: coarsening, neocrystallization, metasomatism, anatexis, and deformation. This aureole contains evidence of these




Figure 4.1 Sample sites (numbered) and metamorphic zones as defined by mineral assemblages and textures. Zones separated by inferred isograds.

processes, locally in combination. Bergantz (1991) defined a metamorphic aureole by changes in texture, composition, and mineralogy. Both definitions are consistent with the contact aureole of the Halifax Pluton.

Based on aforementioned changes in mineralogy and texture, it is possible to infer metamorphic isograds (Fig. 4.1). Evidence for a common reaction in metamorphic aureoles of pelitic rocks (Pattison and Tracy, 1991) is seen from petrographic analysis. Further, an interpretation of the pressure and temperature conditions during formation are inferred from the same information. Microscopic analysis indicates that metamorphism and deformation in the Kearney Lake area is directly related to pluton emplacement.

4.2 Metamorphic Aureole Zonation

Three distinct zones are defined based on mineral assemblages and textures from samples at the northeast end of the Halifax Pluton (Fig. 4.1). The outer zone (biotite zone), is only a small part of the study area, but extends beyond the limit of mapping (Fig. 4.1). The middle zone (cordierite zone), a little less than 750 m wide, extends over the entire length of the northeast end of the Halifax Pluton. The inner zone (K-feldspar zone), is directly adjacent to the contact, approximately 250 m wide, and has a lateral extent similar to the cordierite zone.

4.2.1 Biotite Zone

The biotite zone is interpreted to begin about 1.5 km from the pluton contact (with increasing grade toward the pluton), and represents the transition from regional to contact

metamorphism. Small (<1 mm) biotite porphyroblasts, a fine grained unaltered matrix, a lack of cordierite, and the dominance of phyllosilicate-defined regional S_1 cleavage characterize this zone. The peak mineral assemblage is biotite, chlorite, quartz, and muscovite. The biotite zone extends beyond the limit of sampling, however the boundary with higher grade rocks is defined where cordierite enters the assemblage, at the cordierite-in isograd. A transition zone (Fig. 4.1) is interpreted between the biotite and cordierite zones, here the assemblages and textures represent a combination of both zones.

Sample 13 is typical of this biotite zone (Fig. 4.2). The sample consists of numerous, <1 mm sized, randomly oriented biotite porphyroblasts in a very fine-grained matrix of quartz and muscovite. Muscovite is subtly aligned approximately parallel to the regional axial planar cleavage (S_1), 240° . The sample contains fresh biotite and no indications of deformation effects.

The biotite zone represents both the outer limits of contact metamorphism, and the transition to regional metamorphism. The exact transition surface from biotite zone to regional grade is not defined accurately. Muecke and Keppie (1979) have defined the regional metamorphic grade of the area to be of chlorite/biotite grade.

4.2.2 Cordierite Zone

The presence of cordierite, an increase in muscovite abundance, and grain size coarsening, define the cordierite zone. Chloritized cordierite porphyroblasts ("eyes") are omnipresent, and attain relatively large sizes, >3 mm, giving the slates in this zone a spotted look (samples 1, 8, and 17 are excellent examples). Minor sphene is seen in most samples from this




Figure 4.2 Sample 95-13, typical texture and mineral assemblage of Biotite zone. Dark ovals are biotite, fabric trends parallel to crack in slide. Horizontal section, north right, bar is 2 mm.

zone, forming very tiny, <1 mm well rounded grains. The peak mineral assemblage is cordierite, biotite, muscovite, quartz, and plagioclase. Most porphyroblasts of cordierite are completely chloritized (retrograde), and only its characteristic ovoid crystal habit allows identification. Cordierite and biotite porphyroblasts are randomly oriented, and inclusion trails are not observed (Fig. 4.3). The outer zone boundary is the cordierite-in isograd, and the inner boundary is the muscovite-out isograd. (Fig. 4.1).

The muscovite-defined foliation intensifies with decreasing distance from the contact in this zone. Porphyroblast-rich samples show a foliation-intensification at porphyroblast edges, and between adjacent porphyroblasts (Fig. 4.4). Muscovite in these areas is coarser grained and better developed than elsewhere within the matrix. This well developed muscovite alignment trends approximately parallel to the pluton boundary. Samples further from the contact have foliation trends oblique to the contact and intermediate between S_1 and S_2 (Fig. 3.3).

Grain size in the cordierite zone is larger on average than in the biotite zone. Textures indicate some recrystallization; strained quartz, augen structures, and grain coarsening (e.g. samples 4, 5, 17, and 8). Some lithological differences lead to textural and mineralogical variation of the host rock. For example, sample 21 consists of calc-silicate nodules which contain amphibole and epidote (not seen in any other sections). Similar observations have been interpreted as metasomatic effects (Purves, 1974).

There are noticeable differences in rock mineralogy and texture within the cordierite zone. Variation, related to lithological layering is expected, however, and therefore isograd placement




Figure 4.3 Sample 95-17, representative of Cordierite zone. Randomly oriented, large cordierite porphyroblasts (dark ovals) in fine grained matrix. Fabric running diagonally (lower left corner), north up. Horizontal section, bar is 4mm.




Figure 4.4 Sample 95-8a, illustrating muscovite-defined-cleavage intensification on edges and in between adjacent cordierite porphyroblasts. Oval grains are altered (chloritized) cordierites, light areas in between aligned muscovites, north up. Horizontal section, bar is 1 mm.

was based on a combination of textures and mineral assemblages. Samples closer to the contact show increasing evidence of recrystallization and an intensification of contact parallel cleavage.

4.2.3 *K-feldspar Zone*

The K-feldspar zone contains the highest grade rocks of the aureole, recrystallized massive LS₂ tectonites. The contact-parallel LS₂ fabric is characteristic of this zone, where complete recrystallization is evident from the fabric, unstrained quartz, and an increased grain size (Fig. 4.5).

Intergranular cordierite, K-feldspar, and a lack of muscovite characterize this zone. Muscovite is rare and where present, grains are irregularly shaped, embayed, and ragged. The feldspars are large, and some samples show myrmekite textures. Sphene is present as tiny crystals in some samples. The peak mineral assemblage is K-feldspar, biotite, quartz, plagioclase and cordierite. This zone is approximately 250 meters wide, bounded by the Halifax Pluton and the interpreted muscovite-out isograd, respectively.

Foliations (S₂) are visible in some of the sections (14 and 15 contain the best developed examples). This foliation, while "wavy", is oriented approximately parallel to the contact. Interestingly, S₂ does not appear as well defined in thin section as in outcrop or hand sample scale. S₂ is defined by quartz+K-feldspar-rich zones separated by biotite-rich bands. The biotite-rich bands have weak grain alignment in samples far from the contact (e.g. samples 14 and 15).

Lineations (L₂) in this zone (samples 19, and 19a in particular) are defined by biotite-rich aggregates, with some grain alignment. L₂ shows better grain (biotite) alignment than any of the




Figure 4.5 Sample 95-14, significant grain coarsening and S_2 (subtle at thin section scale) running top left-bottom right, north up. Horizontal section, bar is 2 mm.

other fabrics. Cordierite is altered and strung out between these biotite bands (Fig. 4.6a and b). The biotite-rich bands anastomose and pinch and swell along their length, but indicate no consistent asymmetry (Fig. 4.6a).

Sample 56 illustrates the characteristic mineralogy and textures from the K-feldspar zone particularly well. Biotite, feldspar, and quartz make up the bulk of this sample. The "wavy" foliations described earlier are well developed in this section, however ring-like structures dominate some parts of the sample (Fig. 4.7a). The origin of the "rings" is not clear. The biotite exhibits a deep rich brown color with numerous pleochroic haloes (Fig. 4.7b). Quartz shows no signs of strain, and there is a total lack of muscovite. The feldspars are mostly K-feldspar, orthoclase or possibly sanidine specifically (as defined from 2V angle determination), indicating rapid cooling and therefore shallow level, low pressure environment of formation (pers. comm. Dr. R. Jamieson, 1996). Conspicuous grain boundary zonation, myrmekite alteration (Fig. 4.8), and the proximity of the sample to the contact (Fig. 4.1), indicate a high diffusion rate.

4.3 Similar Contact Aureoles

Metamorphic contact aureoles are common in Meguma-like sedimentary rocks. The Comrie aureole of Scotland (Tilley, 1924; Paterson et al., 1991; Pattison and Tracy, 1991; Paterson and Vernon, 1995), the aureole of the Duluth Complex (Labotka et al., 1981), and the Oslo hornfelses (Goldschmidt, 1911), are perhaps the most analogous, well studied metamorphic contact aureoles. Pattison and Tracy (1991) describe mineralogical, textural, and compositional




Figure 4.6a) Sample 95-19, intergranular cordierite (dark gray masses) strung out between wavy L_2 , notice biotite grain alignment in L_2 bands, north right. Horizontal section, bar is 4 mm.




Figure 4.6b) Sample 95-19a, vertical section showing biotite grain alignment (black) and intergranular cordierite (light gray). Bar is 1 mm.




Figure 4.7a) Sample 95-56, biotite (black) defined ring-like structures. K-feldspar, quartz, and plagioclase within rings. Fabric running top left-bottom right. Horizontal section, bar is 4 mm.




Figure 4.7b) Sample 95-56, randomly oriented biotite grains (dark and elongate) defining the ring-like structures. Horizontal section, bar is 1 mm.




Figure 4.8 Sample 95-56, mymekite alteration of plagioclase (dark gray) by K-feldspar (light gray). Horizontal section, bar is 0.25 mm.


changes within the Comrie aureole comparable to those of Halifax Pluton aureole. Diagnostic features include the presence of cordierite, absence of andalusite, and muscovite-quartz breakdown with associated K-feldspar. The Comrie aureole therefore serves as an excellent model. Pattison and Tracy (1991) interpret the Comrie sequence to represent a low pressure, low temperature environment, and present a sequence of model reactions for the Comrie aureole.

Two of the more pertinent reactions for the Halifax Pluton are:

Cordierite-in isograd (Halifax Pluton aureole)



Muscovite-out isograd (Halifax Pluton aureole)



Comparable reactions appear to operate at both the Comrie aureole and the Halifax Pluton aureole. These two reactions (4.1 and 4.2) seem consistent with the interpreted isograds, indicating the growth of cordierite and the progressive elimination of muscovite, respectively. It appears the Halifax Pluton aureole experienced contact metamorphic conditions similar to the Comrie aureole.

Field descriptions of Pattison and Tracy (1991) indicate a variation in mineralogy and texture with distance from the contact that closely resembles variation in the Halifax Pluton

aureole. These descriptions include a decrease in fissility of hornfels and anatexis at the granite boundary. Similar conditions (e.g. temperature, pressure, and depth of formation) presumably existed during formation of each contact aureole.

4.4 Timing of Contact Metamorphism and Deformation

Within the LS₂ schist zone, metamorphism and deformation appear synchronous. Evidence for this is seen in sample 95-19 (Fig. 4.6a and b). Figure 4.6a illustrates the biotite bands anastomosing around cordierite porphyroblasts. Figure 4.6b shows inclusions of biotite grains within the intergranular cordierite, aligned with the biotite bands. Both observations indicate that L₂ formed when cordierite was growing, suggesting a synchronous relationship between metamorphism and deformation. In addition, high temperature metamorphic minerals K-feldspar, plagioclase, and recrystallized quartz define the LS₂ fabric, the deformation responsible for the fabric, must have occurred with metamorphism.

Further from the contact, metamorphism appears to have been initiated prior to deformation. Porphyroblasts of cordierite grew unaffected by any foliation. However, they deflect a contact-parallel foliation. A thermal pulse preceding the magma plume may be responsible for this initial growth. Post-cordierite strain presumably formed the contact-parallel structures, e.g. LS₂, boudins, the Kearney Lake Anticline, and the muscovite defined foliation of the cordierite zone.

4.5 Conclusions

Petrographic analyses allow the metamorphic contact aureole of the Halifax Pluton to be interpreted in terms of a set of metamorphic zones. Figure 4.9 summarizes all mineral assemblages, and Appendix C tabulates all petrographic observations.

Metamorphism adjacent to the northeast end of the Halifax Pluton is related to pluton emplacement. With this conclusion plus compositional, mineralogical, and textural evidence; pressure and temperature conditions of formation were concluded (Fig. 4.10). Figure 4.10 is a phase diagram indicating under what conditions, inferred reactions may occur. The phase diagram constrains the pressure and temperatures to approximately 2.5 kb and 550-700 °C, respectively. The work done on the similar Comrie aureole corresponded, with interpretations for temperatures of 500-600 °C, at pressures of approximately 3 kb (Pattison and Tracy, 1991).

The interpreted isograds, while roughly positioned, indicate specific changes in mineralogy and textures directly related to contact metamorphism. Both field and microscopic evidence lead to the conclusion that at least a 1.5 kilometer wide zone of the Meguma Group was subjected to contact metamorphism, as a result of Halifax Pluton emplacement.




Figure 4.9 Summary of mineral assemblages. Isograds separate distinct zones. Stippled area represents transition zone.




Figure 4.10 Phase diagram of metamorphic minerals present in K-feldspar zone. Andalusite (And), Sillimanite (Sil), and Kyanite (Ky) field for reference. Line (a) indicates reaction ($\text{Mus} + \text{Bt} + \text{Qtz} \Rightarrow \text{H}_2\text{O} + \text{Kfs} + \text{Crd}$), line (b) ($\text{Mus} + \text{Qtz} \Rightarrow \text{Kfs} + \text{Al}_2\text{SiO}_5 + \text{H}_2\text{O}$), and line (c) ($\text{Kfs} \Rightarrow \text{Sanadine}$). Typical compositions for metamorphic minerals were used in calculations: 80% Mg-Crd and 50% Phlogopite. Phase diagram determined with PTXSS software after Brown et al., 1979.

CHAPTER 5 DISCUSSION AND CONCLUSIONS

5.1 Introduction

Evidence for the mechanism of emplacement surrounding the South Mountain Batholith is rare. Until recently the batholith was considered to have been emplaced by exclusively passive mechanisms. This study and the work of Burns (1995) have led to recognition of emplacement-related structures that suggest a more active mode of emplacement along the northeast flank of the Halifax Pluton. Field mapping for this study documented a metamorphic contact aureole containing several deformation structures previously unknown in this area. Detailed macroscopic and microscopic work has determined that deformation and metamorphism in the contact aureole were essentially synchronous. No overprinting relationship between contact and regional metamorphism was recognized in this study. This new evidence sheds light regarding the mode of emplacement for the Halifax Pluton.

The previous chapters have documented data which defined a metamorphic contact aureole and zone of emplacement-related strain. Interpretations have been left to the present chapter, where constraints on pluton-associated strain on the country rocks will be discussed.

5.2 Cross-section

A cross-section along the NE-SW hinge of the Magazine Hill Basin was presented by Burns (1995). The present study has augmented the cross-section with the addition of S_0 and S_2 data close to the contact (Fig 5.1). S_0 and S_2 measurements lying close to the cross-section line were projected back along strike to the line, where apparent dips were calculated and subsequently drawn on the vertical plane. The moderately dipping beds of the Magazine Hill Basin, and the much more steeply dipping beds of the transverse anticline are evident. Beds dip gently far from the contact, but steepen considerably closer to the contact. A detailed section of the LS_2 schist zone illustrates that LS_2 generally dips more steeply than bedding (S_0) (Fig. 5.1 inset). The cross-section clearly shows a decrease in strain with distance from the contact and therefore reinforces Burn's (1995) conclusion that these features are related to pluton emplacement.

Additionally, the cross-section allows an interpretation of the profile geometry of the transverse anticline. The cross-section (Fig. 5.1) indicates that parallel disharmonic folding is not possible, rather similar folds with limb thinning are more likely adjacent to the Halifax Pluton. Therefore, the transverse fold is probably one of a harmonic stack that may propagate downwards as far as layered rock extends adjacent to the pluton. Significant strain (shortening perpendicular to the contact) would be accommodated in the zone close to the contact according to this model of deformation. Such folding would be expected in rocks adjoining the contact, where the heated rocks may act in a ductile manner. With increasing distance from the pluton-country rock




Figure 5.1 Cross-section from A-F line from Figure 3.2 (drawn parallel to hinge of Bedford Syncline and Magazine Hill Basin). Inset highlights S₀ and S₂ dips within LS₂ schist zone.

contact, temperatures diminish dramatically, and ductile deformation is no longer feasible (Paterson et al., 1991b). There the Magazine Hill Basin is a parallel fold and therefore probably detaches at a relatively shallow depth.

Paterson and Vernon (1995) argue that strains inflicted on country rocks by an expanding pluton must be very high near the contact, but decrease in intensity with distance from the contact, due to the increased volume of rock being deformed. Paterson and Vernon (1995) also conclude that structural (strain) aureoles are much less extensive than thermal aureoles. The Magazine Hill Basin, if it is interpreted as emplacement-related, makes the strain aureole of the Halifax Pluton significantly wider than its thermal aureole. Therefore, the Magazine Hill Basin may warrant further investigation.

5.3 Strain Path at the Granite Contact

In order to estimate the strain path of the marginal schist (LS_2) in the Goldenville Formation, model 2-D strain paths were calculated using MathCAD, and compared with field data. The relevant field observations are as follows: L_2 plunges steeply and represents the maximum finite elongation direction, S_0 and foliation within the LS_2 schist zone (S_2) dip towards the granite contact; S_2 generally dips more steeply than S_0 ; the granite contact is vertical (Fig. 5.2a); a small wavelength re-fold lies just beyond the schist zone; and a large wavelength basin separates the re-fold from any rocks unaffected by pluton intrusion.




Figure 5.2a) Observed geometry of S_0 , S_2 , and vertical granite contact. b) Schematic representation of assumed initial geometry. c) Orientations of S_0 and S_2 produced from calculations of exclusively simple shear. d) Orientations of S_0 and S_2 produced from a high pure shear-simple shear combination. e) Intermediate steps and final orientation of granite dipping S_0 produced from calculations using a large pure shear to simple shear ratio.

Simplifying assumptions were required in order to reproduce, as closely as possible, the observations. These assumptions are: the initial trace (prior to any intrusion-related deformation) of S_0 on the section plane is horizontal or possibly slightly inclined to the southwest, S_2 represents the trace of the XY plane of the strain ellipsoid (as shown by minor structures associated with LS_2) (pers. comm. N. Culshaw, 1996), and simple shear exerted by granite intrusion on the Goldenville would have been granite-side-up (Fig. 5.2b). The assumptions are necessary, but do entail some degree of approximation. For instance, the orientation of the granite contact may not be perfectly vertical. Also the model pure and simple shear strain paths are two dimensional (confined to the section plane). This is a moderate oversimplification in view of the Y-parallel (3 dimensional) extension evident from field observations of the orthorhombic LS_2 fabric (Fig. 3.6a). Using the previous assumptions, model strain paths were calculated. Assuming that the granite boundary is vertical, and exclusively a simple shear plane, incorrect orientations of S_0 and S_2 were produced (Fig. 5.2c). A combination of simple shear and pure shear results in orientations close to the observed. S_0 and S_2 orientations did not exactly match the observed, rather they were dispersed to either side of the shear plane (Fig. 5.2d). What is important, however, is that only a high ratio of pure shear to simple shear produced S_0 that dipped toward the granite. Therefore, model results suggest that pure shear (i.e. expansion of the pluton perpendicular to its boundary) was more important than simple shear in pluton emplacement, and associated deformation of the Meguma Group in the ductile aureole.

The geometry of the transverse fold in the same section plane (Fig. 5.1) is further evidence for flattening perpendicular to the contact. The steep S_0 measurements adjacent to the contact, define an asymmetric fold. Post-emplacment, regional deformation does not seem a reasonable source for its configuration

5.4 Field and Microscopic Interpretations

Microscopic evidence supplements field observations and implies that metamorphism and deformation were synchronous. Mineral assemblages are consistent with well known contact metamorphic reactions, indicating that metamorphism was due to pluton emplacement. Trends of S_1 cleavage swing from regional to contact-parallel (LS_2). This may be interpreted to indicate that S_1 and S_2 are the same structure, with separate orientations caused by deformation of the country rock during pluton emplacement (Guglielmo, 1992). This interpretation would entail syntectonic, or perhaps late syntectonic plutonism (Guglielmo, 1992).

5.5 Halifax Pluton Emplacement

The above observations fit well with current models of rising plutons; a thermal pulse precedes the main bulk of the magma, hot magma is then emplaced in cold country rocks, metamorphism ensues. Subsequently, the tail of the pluton rises, pushing out the now crystallized pluton margins and initiating another stage of deformation (Paterson et al., 1991b). In these models, pluton expansion causes flattening of adjacent country rock, hence schistose foliations

form in both the granite and deformed country rock. Marginal foliations in the granite are not evident in the Halifax Pluton; although a country rock strain aureole does exist, containing foliations (LS_2).

L_2 structures may indicate some vertical movement between the pluton and the country rocks, perhaps emplacement derived, or perhaps from post-emplacement regional deformation. The latter seems unlikely due to the steeply plunging nature of L_2 on the foliation plane. Evidently, L_2 is syn-plutonic and indicates a vertical material transfer was active during pluton emplacement (Paterson et al., 1991b).

Regional deformation can play an additional role in material transfer during pluton emplacement. Paterson and Vernon (1995) argue that regional deformation accompanying emplacement is of only minor importance. Their evidence comes in the form of a lack of far-field (> 0.4 of pluton radius) displacements around the studied plutons (Ardara, Ireland; Cannibal Creek, Australia; Papoose Flat, California). The anomalous Magazine Hill Basin may be an example of such far-field displacement.

5.6 Conclusions

This study recognized and documented some regionally anomalous structures in the vicinity of Kearney Lake, defined a contact aureole, and inferred relationships between metamorphism and deformation. The main results of this investigation indicate contact metamorphism and deformation were intimately related and the emplacement-derived nature of

these local structures as interpreted by Burns (1995) was reinforced. Further work outside the limited study area may answer some of the remaining questions. Specifically, further internal examination Halifax Pluton (in an attempt to define a fabric) and an investigation of southwest and northwest pluton boundaries (to find the LS₂ schist). As well, identification of any further regionally anomalous structures may shed more light on pluton-emplacment processes involved with the Halifax Pluton, and indeed the South Mountain Batholith.

REFERENCES

- Abbott, R.N. 1989. Internal Structures in part of the South Mountain Batholith, Nova Scotia, Canada. *Geological Society of America Bulletin*, **101**, pp.1493-1506.
- Bateman, R. 1985. Aureole Deformation by Flattening Around a Diapir During In Situ Ballooning: The Cannibal Creek Granite. *Journal of Geology*, **93**, pp.293-310.
- Bergantz, G.W. 1991. Chemical and physical characterization of Plutons, in Kerrick, D.M., ed., *Contact Metamorphism: Mineralogical Society of America Reviews in Metamorphism*, **26**, pp.13-42.
- Brown, T.H., Berman, R.G., and Perkins, E.H. 1988. GeO-CALC: Software package for calculation and display of pressure-temperature-composition phase diagram diagrams using an IBM or compatible personal computer. *Computers and Geosciences*, **14**, pp.279-289.
- Burns, N.R. 1995. Structural Geology of the Meguma Group adjacent to the eastern contact of the South Mountain Batholith. B.Sc. Honors thesis, Dalhousie University.
- Clarke, D.B., and Chatterjee, A.K. 1988. Physical and Chemical Processes in the South Mountain Batholith. Recent Advances in Geology of Granite Related Mineral Deposits. In: Special Volume, Canadian Institute of Mining and Metallurgy, pp.223-233.
- Clarke, D.B., Chatterjee, A.K., and Giles, P.S. 1993. Petrochemistry, tectonic history, and Sr-Nd systematics of the Liscomb Complex, Meguma Lithotectonic Zone, Nova Scotia. *Canadian Journal of Earth Sciences*, **30**, pp.449-464.
- Clarke, D.B. and Tate, M.C. 1993. Origin of late Devonian Weekend lamprophyre dykes Meguma Zone, Nova Scotia. *Canadian Journal of Earth Sciences*, **30**, pp.2295-2305.
- Faribault, E.R. 1908. City of Halifax sheet Map No. 68; Geological Survey of Canada; publication No. 44.

- Fyson, W.K. 1980. Fold Fabrics and emplacement of an Archean granitoid pluton, Cleft Lake, Northwest Territories. *Canadian Journal of Earth Sciences*, **17**, pp.325-332.
- Goldschmidt, V.M. 1911. Die Kontakt Metamorphose in Kristianiagebiet. *Lkr. Videnskabselsk, Kristiana* 1, 483 p.
- Guglielmo, G. Jr. 1992. Magmatic strains and Foliation triple points of the Merrimac plutons, northern Sierra Nevada, California: implications for pluton emplacement and timing of subduction. *Journal of Structural Geology*, **15**, pp.177-189.
- Guglielmo, G. Jr. 1994. Interference between pluton expansion and coaxial tectonic deformation: three dimensional computer model and field implications. *Journal of Structural Geology*, **16**, No.2, pp.237-252.
- Henderson, J.R., Wright, T.O., and Henderson, M.N., 1986. A history of cleavage and folding: An example from the Goldenville Formation, Nova Scotia. *Geological Society of America Bulletin*, **97**, pp.1354-1366.
- Horne, R.H. and Culshaw, N. 1993. Preliminary Evaluation of Flexural Slip and its significance in localizing auriferous veins in the Meguma Group, Nova Scotia. Nova Scotia Department of Natural Resources, Mines and Energy Branches Report 94-1, pp.147-160.
- Horne, R.H., MacDonald, M.A., Coney, M.C., and Ham, L.J. 1992. Structure and emplacement of the South Mountain Batholith, southwestern Nova Scotia. *Atlantic Geology*, **28**, pp.29-50.
- Keppie, J.D. 1982. The Minas Geofracture. Major Structural Zones and Faults of the Northern Appalachians. Edited by P.St-Julien and J. Beland. *Geological Association of Canada, Special Paper*, **24**, pp.263-280.
- Keppie, J.D., and Dallmeyer, R.D. 1987. Dating transcurrent terrane accretion: an example from the Meguma and Avalon composite terranes in the northern Appalachians. *Tectonics*, **6**, pp. 831-847.

- Kerrick, D.M., 1991. Contact Metamorphism, an introduction. In: Contact Metamorphism, Mineralogical Society of America Reviews in Mineralogy, **26**, pp.1-13.
- Kontak, D.J., Cormier, R.F., Reynolds, P.H., and Taylor, K. 1989. Preliminary results of Rb/Sr and $^{40}\text{Ar}/^{39}\text{Ar}$ geochronological investigation, East Kemptville leucogranite, southwestern Nova Scotia: Evidence for a ca 370 Ma age of emplacement and multiple tectono-thermal overprinting events. Mines and Minerals Branch Report of Activities 1989, Part A. Edited by D.R. MacDonald and K.A. Mills. Nova Scotia Department of Mines Report, 89-3, pp.41-47.
- Labotka, T.C., Papike, J.J., Vaniman, D.T., and Morey G.B. 1981. Petrology of contact metamorphosed argillite from the Rove Formation, Gunflint Trail, Minnesota. *American Mineralogist*, **66**, pp.70-86.
- Lajtai, E.Z., and Stringer, P. 1981. Joints tensile strength and preferred fracture orientation in sandstones, New Brunswick and Prince Edward Island, Canada. *Maritime Sediments and Atlantic Geology*, **17**, pp.70-87.
- Muecke, G. and Keppie, J.D. 1979. Metamorphic geology map of Nova Scotia. Nova Scotia Department of Mines and Energy.
- MacDonald, M.A., Horne, R.J., Corey, M.C., and Ham, L.J. 1992. An overview of recent bedrock mapping and follow-up petrological studies of the South Mountain Batholith, southwestern Nova Scotia, Canada. *Atlantic Geology*, **28**, pp.7-28.
- Paterson, S.R., Brudos, T., Fowler, K., Carlson, C., Bishop, K., and Vernon, R.H. 1991a. Papoose Flat pluton: Forceful expansion or postemplacement deformation? *Geology*, **19**, pp.324-327.
- Paterson, S.R. and Fowler, T.K. Jr. 1993. Re-examining pluton emplacement processes. *Journal of Structural Geology*, **15**, pp.191-206.
- Paterson, S.R. and Vernon, R.H. 1995. Bursting the bubble of ballooning plutons, A return to nested diapirs emplaced by multiple processes. *Geological Society of America Bulletin*, **107**, pp.1356-1380.

- Paterson, S.R., Vernon, R.H., and Fowler, T.K. Jr. 1991b. Aureole tectonics. In: Contact Metamorphism. Mineral Society of America Reviews in Mineralogy, **26**, pp.673-722.
- Pattison, D.R.M. and Tracy R.J., 1991. Phase Equilibria and Thermobarometry of Metapelites. In: Contact Metamorphism. Mineral Society of America Review in Mineralogy, **26**, pp.105-205.
- Purves, M.F. 1974. Metasomatism of calcareous concretions of the Goldenville Formation, Nova Scotia. B.Sc. Honors thesis, Dalhousie University.
- Schenk, P.E. 1981. The Meguma Terrane of Nova Scotia, Canada- An aid in trans-Atlantic correlation, In: Schenk, P.E., ed., Regional trends in the geology of the Appalachain-Caledonian-Hercynian-Mauritanide orogen: D. Reidel Publishing Company pp.121-130.
- Schenk, P.E. 1991. Events and sea level changes on Gondwana's margin: The Meguma Zone (Cambrian-Devonian) of Nova Scotia Canada. Geological Society of America Bulletin, **103**, pp. 512-521.
- Tilley, C.E. 1924. Conatact-metamorphism in the Comrie area of the Perthshire highlands. Quaternary Journal of the Geological Society of London, **80**, pp.20-71.
- Williams, H. 1979. Appalachian orogen in Canada. Canadian Journal of Earth Sciences, **16**, pp.792-807.
- Woodcock, N.H. and Underhill, J.R. 1987. Emplacement-related fault patterns around the Northern Granite Arran, Scotland. Geological Society of America Bulletin, **98**, pp.515-527.
- Yardley, B.D. 1989. An introduction to Metamorphic Petrology. Longman Scientific & Technical, New York.

Appendix A: Tabulated Field Data

STATION #	EASTING	NORTHING	BEDDING		CLEAVAGE (S_1) FOLIATION (S_2)		LINEATIONS (L_2)	
			STRIKE°	DIP°	STRIKE°	DIP°	TREND°	PLUNGE°
PDG-001	444675	4949435	295	28	122	76	S_1	
PDG-002	443925	4950003	310	25				
PDG-003	442864	4949925	249	20				
PDG-004	443755	4949742	330	20				
PDG-005	443703	4949581	105	22	108	73	S_1	
PDG-007	442846	4951132	280	20				
PDG-008	442355	4951100	275	22				
PDG-009	445632	4948210	160	26				
PDG-010	445513	4948193	140	36				
PDG-011	445235	4948752	155	18				
PDG-013	445912	4951698	022	22	032	72	S_1	
PDG-014	439968	4951139			110	70	S_2	22 70
PDG-015	440167	4951185	172	12	114	88	S_2	348 88
PDG-016	440459	4951685			80	82	S_1	
PDG-017	440657	4952252	245	52	247	82	S_1	335 68
PDG-018	440414	4950913			110	80	S_2	12 82
PDG-019	440877	4950902			118	90	S_2	210 89
PDG-020	441140	4950686			120	85	S_2	25 90
PDG-021	441543	4951960	250	53				
PDG-022	445213	4952385	350	17	055	55	S_1	
PDG-023	443468	4952356	246	12	056	82	S_1	
PDG-026	441516	4950397			145	72	S_2	225 74
PDG-028	441913	4950220			148	68	S_2	
PDG-030	442269	4949906			100	90	S_2	
PDG-031	442435	4949766			120	80	S_2	63 80
PDG-034	442896	4949457			110	80	S_2	225 83
PDG-035	443056	4949479			133	72	S_2	230 70
PDG-037	443565	4948964			145	70	S_2	
PDG-038	443585	4949147	135	42				
PDG-039	443598	4949161	135	42				
PDG-040	443081	4949612			130	75	S_2	
PDG-043	444635	4948511	143	38				
PDG-044	444495	4948712			140	70	S_2	
PDG-045	443814	4949173	120	35				
PDG-047	443877	4948974			119	60	S_2	
PDG-048	443878	4948921			120	70	S_2	
PDG-050	443854	4948838			123	68	S_2	230 45 130 18 L_3
PDG-051	443929	4948831			110	58	S_2	238 60
PDG-052	443832	4948772			118	80	S_2	

PDG-053	443817	4948750	340 54			
			112 74			
PDG-056	443578	4948835		118 70	S ₂	
PDG-057	446293	4944711	100 65	88 65	S ₂	220 44
				238 75	S ₂	278 60
PDG-059	446370	4945322		160 85	S ₂	207 74
PDG-060	446332	4945481	126 76			
PDG-061	446079	4945550	105 65	117 68	S ₂	
PDG-062	445798	4946256		325 80	S ₂	
PDG-063	445723	4946435	125 60			225 60
PDG-064	446124	4946313	192 55			270 58
PDG-065	446068	4946825	130 63			227 65
PDG-066	446123	4945942	50 86	230 88	S ₂	230 10
PDG-068	446910	4943547		65 80	S ₂	67 15
PDG-069	446665	4944021		120 80	S ₂	
PDG-070	447246	4943266		275 80	S ₂	
PDG-071	446722	4944037		70 90	S ₂	
PDG-072	446585	4944383		174 81	S ₂	
PDG-075	440757	4950791		122 76	S ₂	
PDG-077	440589	4940932	100 77	120 86	S ₂	
PDG-078	440412	4950573		110 80	S ₂	

* UTM coordinates determined from GPS (Trimble Pathfinder+). Data points differentially corrected with a Bangor Maine, U.S.A. base station. Positions accurate to <5 meters.

APPENDIX B: LS₂ Schist Zone Measurements

LS₂ foliation measurements (strike and dip in degrees)

124°/89°
119°/85°
120°/86°
122°/88°
121°/89°
120°/80°

S₀ measurements (strike and dip in degrees)

108°/78°
100°/77°
099°/80°
102°/80°

Shear Band-like planar feature measurements (strike and dip in degrees)

287°/51°	280°/52°
290°/55°	282°/48°
287°/62°	283°/45°
276°/50°	279°/50°
281°/63°	283°/42°
285°/65°	285°/48°
282°/53°	278°/54°
284°/54°	282°/52°
280°/35°	278°/47°
284°/50°	280°/53°
281°/38°	283°/59°
281°/38°	283°/59°
281°/48°	288°/52°

APPENDIX C: TABLE OF PETROGRAPHIC DESCRIPTIONS

SAMPLE #	MINERAL ASSEMBLAGE	TEXTURE	PORPHYROBLASTS	ALTERATION	COMMENTS (Distance from contact)
95-13	Quartz, Muscovite, Biotite, Chlorite, and Sphene	Very fine Grained Cleavage -Mus. defined	Small (<1mm), randomly oriented Biotite	Unaltered	Weakly defined foliation wrap around P'blasts. Shadow zones. Rounded Biotites. (3350 meters)
95-23	Quartz, Biotite, opaques, Muscovite, chlorite	Fine Grained Cleavage -Mus. Defined Bedding compositional	Biotite	Some Biotite is chloritized	Bedding~240° Cleavage~068° Irregular Biotite Quartz vein (2510 meters)
95-21	Quartz, Amphibole, Chlorite, Sphene, Biotite	Medium Grained. Large Quartz Grains. Bedding -compositional	Acicular amphibole		Bedding~245° Amphibole alignment? Well preserved clastic texture. Calc-silicate 1550 meters
95-7	Cordierite, Biotite, Quartz, Muscovite, Sphene	Fine Grained. Cleavage -Mus. Defined	Cordierite. Biotite Magnetite Many P'blasts	Chloritization of Cordierite	Opagues equant and omnipresent. Quartz vein. Cleavage~255° (1400 meters)
95-17	Quartz, Biotite, Muscovite, Opaques, Cordierite, some Plagioclase.	Medium Grained. Strong Cleavage -Mus. defined Bedding -compositional -grain sizes differences	Large Cordierite. Red-brown Biotite.	Chloritization of micas, and minor cordierites. Biotite totally replaced by chlorite.	Enhanced Fabric development at P'blast sides. Major Augen wrap around structures. Bedding=Cleavage~245° -Intermediate trend Grain coarsening. Change in Biotite color. (1260 meters)
95-8 95-8a	Quartz, Muscovite, Biotite, Chlorite, Cordierite, opaques (magnetite)	Fine grained. Cleavage -Mus/Bt defined	Cordierites Large (~3-4 mm)	Chloritization of Cordierites - rimmed alteration Biotites altered to chlorite	Fabric enhanced near and between P'blasts. Augen wrap around structures. Cleavage~215° Highly altered-retrograde (1170 meters)
95-1	Quartz, Muscovite, Biotite, Cordierite, opaques, Chlorite	Fine Grained Cleavage -Mus defined	Cordierites (~3mm) Biotites (>1mm)	Chloritization of Cordierites and Biotites. Retrograde chlorite.	Fabric enhanced at P'blast edges. Cleavage~285° Alteration evident (800 meters)

95-4	Quartz, Feldspar, Muscovite, Biotite, Chlorite, Sphene, Hematite.	Medium Grained, clastic texture. Bedding -Compositional Cleavage -Mus/Bt defined	None	Not highly Chlorite replacing Bt	Chlorite well developed Bedding~125° Recrystallized No Cordierites-- lithological? Weak foliation. (860 meters)
95-5	Quartz, Feldspar, Muscovite, Biotite, Cordierite, Chlorite, Sphene, Garnet	Medium Grained. Compositional, pelitic and clastic layers. Bedding? Cleavage -Mus defined	Biotite Cordierite concn. in pelitic layers	Chloritization of Biotite and Cordierite.	Intense fabric around P'blasts. -P'blasts appear rotated -Cleavage anastomoses Cleavage~125°--parallel to bedding. Compositional layering. Intense retrogression (630 meters)
95-47	Plagioclase, Biotite, Quartz, Amphibole, Sphene, Muscovite, opaques, Hematite	Medium Grained. feldspar/quartz coarsening.	Muscovite. Oxides.	Muscovite highly altered. Feldspar altered to Clinzoisite and chlorite	Clastic texture Recrystallized (420 meters)
95-49	Plagioclase, Biotite, Cordierite, Muscovite, opaques	Medium grained.	Corierite	Muscovite beginning to alter on edges	Recrystallized. foliation~100° (350 meters)
95-19	Quartz, Feldspar, Plagioclase, Biotite, Cordierite	Medium Grained. Lineation -Bt alignment in zones	Cordierite granular and strung out and aligned in L ₂	Cordierites reacting out.	L ₂ Biotite defined lineation pinch and swell structure. Minor Mus. -replacement (261 meters)
95-15	Quartz, Plagioclase, Biotite, Cordierite, Muscovite, Sphene, opaques	Medium Grained. Gneissic-recrystallized quartz rich Foliation -Bt/Mus Defined	Cordierite but pretty grungey	Cordierite Plagioclase -Sericite	Cordierite and Biotite well developed. Foliation~120° Cordierite granular Partly preseved clastic texture (110 meters)
95-51	Quartz, Biotite, Muscovite, Feldspar, Plagioclase	Medium to fine grained. Gneissic-recrystallized	None	Muscovite altered, and reacting out	Foliation defined by linear biotite zones, and grain sizes. (150 meters)
95-14	Quartz, Feldspar, Plagioclase, Muscovite, Biotite -both retrograde	Coarse Grained. Recrystallized Foliation -Banding of Bt/Qtz	None	Symplectite feldspar alteration. Muscovite retrograde Chlorite retrograde	Foliation segregated into bands, wavy development ~120° Quartz-rich lithology (60 meters)

95-56	Quartz, Feldspar, Plagioclase, Biotite, Zircon	Fine to Medium Grained. recrystallized/ annealed.	None	Myrmekite Feldspar alteration	Ring like Structures Granoblastic texture Lack of Muscovite. Grain Boundary zonation. (5 meters)
95-58	Quartz, Biotite, Muscovite, Feldspar, Plagioclase, Zircon	Fine Grained. Massive and recrystallized	Rare	Chloritized Biotite rims, and some completely replced.	Weakly defined foliation -perhaps 2 or 3 directions Rare, but present Muscovite. Around corner of pluton near Burnside Industrial Park. Does not enter into metamorphic zonation